

Environmental Protection Agency

§ 180.960

Chemical	CAS No.
Potassium chloride	7447-40-7
2-Propanol (isopropyl alcohol)	67-63-0
Red cabbage color, expressed from edible red cabbage heads via a pressing process using only acidified water	None
Silica, amorphous, fumed (crystalline free)	112945-52-5
Silica, amorphous, precipitated and gel	7699-41-4
Silica gel	63231-67-4
Silica gel, precipitated, crystalline-free	112926-00-8
Silica, hydrate	10279-57-9
Silica, vitreous	60676-86-0
Soap (The water soluble sodium or potassium salts of fatty acids produced by either the saponification of fats and oils, or the neutralization of fatty acid)	None
Sorbic acid, potassium salt	24634-61-5
Soapbark (Quillaja saponin)	1393-03-9
Sodium alginate	9005-38-3
Sodium chloride	7647-14-5
Syrups, hydrolyzed starch, hydrogenated	68425-17-2
Ultramarine blue (C.I. Pigment Blue 29)	57455-37-5
Urea	57-13-6
Vanillin	121-33-5
Xanthan gum	11138-66-2

[67 FR 36537, May 24, 2002]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 180.950, see the List of CFR Sections Affected, which appears in the Finding Aids section of the printed volume and on GPO Access.

§ 180.960 Polymers; exemptions from the requirement of a tolerance.

Residues resulting from the use of the following substances, that meet the definition of a polymer and the criteria specified for defining a low-risk polymer in 40 CFR 723.250, as an inert ingredient in a pesticide chemical formulation, including antimicrobial pesticide chemical formulations, are exempted from the requirement of a tolerance under FFDC section 408, if such use is in accordance with good agricultural or manufacturing practices.

Polymer	CAS No.
Acetic acid ethenyl ester, polymer with ethenol and (α)-2-propenyl-(ω)-hydroxypoly (oxy-1,2-ethanediy) minimum number average molecular weight (in amu), 15,000	137091-12-4
Acetic acid ethenyl ester, polymer with 1-ethenyl-2-pyrrolidinone	25086-89-9
Acetic acid ethenyl ester, polymer with sodium 2-methyl-2-[(1-oxo-2-propen-1-yl)amino]-1-propanesulfonate (1:1), hydrolyzed, minimum number average molecular weight (in amu), 61,000	924892-37-5
Acrylic acid, polymerized, and its ethyl and methyl esters	None

Polymer	CAS No.
Acrylic acid-sodium acrylate-sodium-2-methylpropanesulfonate copolymer, minimum average molecular weight (in amu), 4,500	97953-25-8
Acrylic acid-stearyl methacrylate copolymer, minimum number average molecular weight (in amu), 2,500	27756-15-6
Acrylic acid, styrene, α-methyl styrene copolymer, ammonium salt, minimum number average molecular weight (in amu), 1,250	89678-90-0
Acrylic acid terpolymer, partial sodium salt, minimum number average molecular weight (in amu), 2,400	151006-66-5
Acrylic polymers composed of one or more of the following monomers: Acrylic acid, methyl acrylate, ethyl acrylate, butyl acrylate, hydroxyethyl acrylate, hydroxypropyl acrylate, hydroxybutyl acrylate, carboxyethyl acrylate, methacrylic acid, methyl methacrylate, ethyl methacrylate, butyl methacrylate, isobutyl methacrylate, hydroxyethyl methacrylate, hydroxypropyl methacrylate, hydroxybutyl methacrylate, lauryl methacrylate, and stearyl methacrylate; with none and/or one or more of the following monomers: Acrylamide, N-methyl acrylamide, N,N-dimethyl acrylamide, N-octylacrylamide, maleic anhydride, maleic acid, monoethyl maleate, diethyl maleate, monoethyl maleate, dioctyl maleate; and their corresponding sodium, potassium, ammonium, isopropylamine, triethylamine, monoethanolamine, and/or triethanolamine salts; the resulting polymer having a minimum number average molecular weight (in amu), 1,200	None
Acrylonitrile-butadiene copolymer conforming to 21 CFR 180.22, minimum average molecular weight (in amu), 1,000	9003-18-3
Acrylonitrile-styrene-hydroxypropyl methacrylate copolymer, minimum number average molecular weight (in amu), 447,000	None
α-alkyl (C ₁₂ -C ₁₅) - ω-hydroxypoly(oxypropylene)poly(oxyethylene)copolymers (where the poly(oxypropylene) content is 3-60 moles and the poly(oxyethylene) content is 5-80 moles), the resulting ethoxylated propoxylated (C ₁₂ -C ₁₅) alcohols having a minimum molecular weight (in amu), 1,500	68551-13-3
Alkyl (C ₁₂ -C ₂₀) methacrylate-methacrylic acid copolymer, minimum molecular weight (in amu), 11,900	None
2H-Azepin-2-one, 1-ethenylhexahydro-, homopolymer	25189-83-7
1,3 Benzene dicarboxylic acid, 5-sulfo-, 1,3-dimethyl ester, sodium salt, polymer with 1,3-benzene dicarboxylic acid, 1,4-benzene dicarboxylic acid, dimethyl 1,4-benzene dicarboxylate and 1,2-ethanediol, minimum number average molecular weight (in amu), 2,580	212842-88-1

§ 180.960

40 CFR Ch. I (7-1-09 Edition)

Polymer	CAS No.
3,5-Bis(6-isocyanatohexyl)-2H-1,3,5-oxadiazine-2,4,6-(3H,5H)-trione, polymer with diethylenetriamine, minimum number average molecular weight (in amu), 1,000,000	87823-33-4
Butadiene-styrene copolymer	None
1,4-Butanediol-methylenebis(4-phenylisocyanate)-poly(tetramethylene glycol) copolymer, minimum molecular weight (in amu) 158,000	9018-04-6
Butene, homopolymer	9003-29-6
2-butenedioic acid (Z)-, monobutyl ester, polymer with methoxyethene, sodium salt, minimum number average molecular weight (in amu), 18,200	205193-99-3
2-Butenedioic acid (Z)-, polymer with ethenol and ethenyl acetate, sodium salt, minimum number average molecular weight (in amu), 75,000	139871-83-3
Butyl acrylate-vinyl acetate-acrylic acid copolymer, minimum number average molecular weight (in amu), 18,000	65405-40-5
Castor oil, ethoxylated, oleate, minimum number average molecular weight (in amu) 2,000	220037-02-5
Castor oil, polyoxyethylated; the poly(oxyethylene) content averages 5-54 moles	None
Chlorinated polyethylene	64754-90-1
Cross-linked nylon-type polymer formed by the reaction of a mixture of sebacoyl chloride and polymethylene polyphenylisocyanate with a mixture of ethylenediamine and diethylenetriamine	None
Cross-linked polyurea-type encapsulating polymer	None
Dimethylpolysiloxane minimum number average molecular weight (in amu), 6,800	63148-62-9
Dimethyl silicone polymer with silica, minimum number average molecular weight (in amu), 1,100,000	67762-90-7
α -(o,p-Dinonylphenyl)- ω -hydroxypoly(oxyethylene) produced by condensation of 1 mole of dinonylphenol (nonyl group is a propylene trimer isomer) with an average of 140-160 moles of ethylene oxide	9014-93-1
Docosyl methacrylate-acrylic acid copolymer, or docosyl methacrylate-octadecyl methacrylate-acrylic acid copolymer, minimum number average molecular weight (in amu), 3,000	None
1,12-Dodecanediol dimethacrylate polymer, minimum molecular weight (in amu), 100,000	None

Polymer	CAS No.
α -(p-Dodecylphenyl)- ω -hydroxypoly(oxyethylene) produced by the condensation of 1 mole of dodecylphenol (dodecyl group is a propylene tetramer isomer) with an average of 30-70 moles of ethylene oxide	9014-92-0 26401-47-8
1, 2-Ethanediamine, polymer with methyl oxirane and oxirane, minimum number average molecular weight (in amu), 1,100	26316-40-5
Ethylene glycol dimethacrylate-lauryl methacrylate copolymer, minimum molecular weight (in amu), 100,000	None
Ethylene glycol dimethacrylate polymer, minimum molecular weight (in amu), 100,000	None
Formaldehyde, polymer with α -[bis(1-phenylethyl)phenyl]- ω -hydroxypoly(oxy-1,2-ethanediyl), number average molecular weight (in amu), 1,803	157291-93-5
Formaldehyde, polymer with 2-methyloxirane and 4-nonylphenol, minimum number average molecular weight (in amu), 4,000	37523-33-4
Fumaric acid-isophthalic acid-styrene-ethylene/propylene glycol copolymer, minimum average molecular weight (in amu), 1×10^{18}	None
Hexadecyl acrylate-acrylic acid copolymer, hexadecyl acrylate-butyl acrylate-acrylic acid copolymer, or hexadecyl acrylate-dodecyl acrylate-acrylic acid copolymer, minimum number average molecular weight (in amu), 3,000	None
Hexamethyl disilazane, reaction product with silica, minimum number average molecular weight (in amu), 645,000	68909-20-6
1,6-Hexanediol dimethacrylate polymer, minimum molecular weight (in amu), 100,000	None
α -Hydro- ω -hydroxy-poly(oxyethylene) C8 alkyl ether citrates, poly(oxyethylene) content is 4-12 moles, minimum number average molecular weight (in amu) 1,300.	330977-00-9
α -Hydro- ω -hydroxy-poly(oxyethylene) C10-C16-alkyl ether citrates, poly(oxyethylene) content is 4-12 moles, minimum number average molecular weight (in amu) 1,100.	330985-58-5
α -Hydro- ω -hydroxy-poly(oxyethylene) C16-C18-alkyl ether citrates, poly(oxyethylene) content is 4-12 moles, minimum number average molecular weight (in amu) 1,300.	330985-61-0
α -Hydro- ω -hydroxypoly(oxyethylene), minimum molecular weight (in amu), 100,000	None
α -Hydro- ω -hydroxypoly(oxyethylene)poly(oxypropylene) poly(oxyethylene) block copolymer; the minimum poly(oxypropylene) content is 27 moles and the minimum molecular weight (in amu) is 1,900	None
α -Hydro- ω -hydroxypoly(oxypropylene); minimum molecular weight (in amu) 2,000	None

Environmental Protection Agency

§ 180.960

Polymer	CAS No.
12-Hydroxystearic acid-polyethylene glycol copolymer, minimum number average molecular weight (in amu), 3,690	70142-34-6
Isodecyl alcohol ethoxylated (2-8 moles) polymer with chloromethyl oxirane, minimum number average molecular weight (in amu) 2,500	None
Lauryl methacrylate-1,6-hexanediol dimethacrylate copolymer, minimum molecular weight (in amu), 100,000	None
Maleic acid-butadiene copolymer	None
Maleic acid monobutyl ester-vinyl methyl ether copolymer, minimum average molecular weight (in amu), 52,000	25119-68-0
Maleic acid monoethyl ester-vinyl methyl ether copolymer, minimum average molecular weight (in amu), 46,000	25087-06-3
Maleic acid monoisopropyl ester-vinyl methyl ether copolymer, minimum average molecular weight (in amu), 49,000	31307-95-6
Maleic anhydride-diisobutylene copolymer, sodium salt, minimum number average molecular weight (in amu) 5,0007-18,000	37199-81-8
Maleic anhydride-methylstyrene copolymer sodium salt, minimum number average molecular weight (in amu), 15,000	60092-15-1
Maleic anhydride-methyl vinyl ether, copolymer, average molecular weight (in amu), 250,000	None
Methacrylic acid-methyl methacrylate-polyethylene glycol methyl ether methacrylate copolymer, minimum number average molecular weight (in amu), 3,700	100934-04-1
Methacrylic copolymer, minimum number average molecular weight (in amu), 15,000	63150-03-8
Methyl methacrylate-methacrylic acid-monomethoxypolyethylene glycol methacrylate copolymer, minimum number average molecular weight (in amu), 2,730	119724-54-8
Methyl methacrylate-2-sulfoethyl methacrylate-dimethylaminoethylmethacrylate-glycidyl methacrylate-styrene-2-ethylhexyl acrylate graft copolymer, minimum average molecular weight (in amu), 9,600	None
Methyl vinyl ether-maleic acid copolymer, minimum number average molecular weight (in amu), 75,000	25153-40-6
Methyl vinyl ether-maleic acid copolymer, calcium sodium salt, minimum number average molecular weight (in amu), 900,000	62386-95-2
Monophosphate ester of the block copolymer α -hydro- ω -hydroxypoly(oxyethylene) poly(oxypropylene) poly(oxyethylene); the poly(oxypropylene) content averages 37-41 moles, average molecular weight (in amu), 8,000	None

Polymer	CAS No.
α -(p-Nonylphenyl)- ω -hydroxypoly(oxyethylene) mixture of dihydrogen phosphate and monohydrogen phosphate esters and the corresponding ammonium, calcium, magnesium, monoethanolamine, potassium, sodium, and zinc salts of the phosphate esters; the nonyl group is a propylene trimer isomer and the poly(oxyethylene) content averages 30 moles	None
α -(p-Nonylphenyl)- ω -hydroxypoly(oxyethylene) sulfate, and its ammonium, calcium, magnesium, monoethanolamine, potassium, sodium, and zinc salts; the nonyl group is a propylene trimer isomer and the poly(oxyethylene) content averages 30-90 moles of ethylene oxide	None
α -(p-Nonylphenyl)- ω -hydroxypoly(oxypropylene) block polymer with poly(oxyethylene); polyoxypropylene content of 10-60 moles; polyoxyethylene content of 10-80 moles; molecular weight (in amu), 1,200-7,100.	None
α -(p-Nonylphenyl)poly(oxypropylene) block polymer with poly(oxyethylene); polyoxyethylene content 30 to 90 moles; molecular weight (in amu) averages 3,000	None
Octadecanoic acid, 12-hydroxy-, homopolymer, octadecanoate minimum number average molecular weight (in amu), 1,370	58128-22-6,
α -cis-9-Octadecenyl- ω -hydroxypoly(oxyethylene); the octadecenyl group is derived from oleyl alcohol and the poly(oxyethylene) content averages 20 moles	None
Octadecyl acrylate-acrylic acid copolymer, octadecyl acrylate-dodecyl acrylate-acrylic acid copolymer, octadecyl methacrylate-butyl acrylate-acrylic acid copolymer, octadecyl methacrylate-hexyl acrylate-acrylic acid copolymer, octadecyl methacrylate-dodecyl acrylate-acrylic acid copolymer, or octadecyl methacrylate-dodecyl methacrylate-acrylic acid copolymer, minimum number average molecular weight (in amu) 3,000	None
Oleic acid diester of α -hydro- ω -hydroxypoly(oxyethylene); the poly(oxyethylene), average molecular weight (in amu), 2,300	None
2-oxepanone, homopolymer, minimum number average molecular weight (in amu) 52,000	24980-41-4
Oxirane, decyl-, reaction products with polyethylene-polypropylene glycol ether with trimethylolpropane (3:1)	903890-89-1
Oxirane, hexadecyl-, reaction products with polyethylene-polypropylene glycol ether with trimethylolpropane (3:1)	893427-80-0
Oxirane, methyl-, polymer with oxirane, ether with 2-ethyl-2-(hydroxymethyl) - 1,3 - propanediol (3:1), reaction products with tetradecyloxirane	903890-90-4

§ 180.960

40 CFR Ch. I (7-1-09 Edition)

Polymer	CAS No.
Oxirane, methyl-, polymer with oxirane, mono[2-(2-butoxyethoxy) ethyl] ether, minimum number average molecular weight (in amu), 2,500	85637-75-8
Oxirane, methyl-, polymer with Oxirane, Monobutyl Ether	9038-95-3
Oxirane, 2-methyl-, polymer with oxirane, minimum number average molecular weight (in amu), 1,100	9003-11-6
Oxirane, 2-methyl-, polymer with oxirane, mono [2-[2-(2-butoxymethylethoxy)methylethoxy]methylethyl] ether, minimum number average molecular weight (in amu), 3,000	926031-36-9
Polyamide polymer derived from sebacic acid, vegetable oil acids with or without dimerization, terephthalic acid and/or ethylenediamine	None
Polyethylene glycol-polyisobutenyl anhydride-tall oil fatty acid copolymer, minimum number average molecular weight (in amu), 2,960	68650-28-2
Polyethylene, oxidized, minimum number average molecular weight (in amu), 1,200	None
Polymethylene polyphenylisocyanate, polymer with ethylene diamine, diethylene triamine and sebacyl chloride, cross-linked; minimum number average molecular weight (in amu), 100,000	None
Polyoxyethylated primary amine (C ₁₄ -C ₁₈); the fatty amine is derived from an animal source and contains 3% water; the poly(oxyethylene) content averages 20 moles	None
Polyoxyethylated sorbitol fatty acid esters; the polyoxyethylated sorbitol solution containing 15% water is reacted with fatty acids limited to C ₁₂ , C ₁₄ , C ₁₆ , and C ₁₈ , containing minor amounts of associated fatty acids; the poly(oxyethylene) content averages 30 moles.	None
Polyoxyethylated sorbitol fatty acid esters; the sorbitol solution containing up to 15% water is reacted with 20-50 moles of ethylene oxide and aliphatic alkanolic and/or alkenolic fatty acids C ₈ through C ₂₂ with minor amounts of associated fatty acids; the resulting polyoxyethylene sorbitol ester having a minimum molecular weight (in amu), 1,300	None
Poly(oxyethylene/oxypropylene) monoalkyl (C ₆ -C ₁₀) ether sodium fumarate adduct, minimum number average molecular weight (in amu), 1,900	102900-02-7
Polyoxymethylene copolymer, minimum number average molecular weight (in amu), 15,000	None
Poly(oxypropylene) block polymer with poly(oxyethylene), molecular weight (in amu), 1,800-16,000	None
Poly(phenylhexylurea), cross-linked, minimum average molecular weight (in amu), 36,000	None

Polymer	CAS No.
Polypropylene	9003-07-0
Polystyrene, minimum number average molecular weight (in amu), 50,000	9003-53-6
Polytetrafluoroethylene	9002-84-0
Polyvinyl acetate, copolymer with maleic anhydride, partially hydrolyzed, sodium salt, minimum number average molecular weight (in amu), 53,000	None
Polyvinylpyrrolidone butylated polymer, minimum number average molecular weight (in amu), 9,500	26160-96-3
Polyvinyl acetate, minimum molecular weight (in amu), 2,000	None
Polyvinyl acetate-polyvinyl alcohol copolymer, minimum number average molecular weight (in amu), 50,000	25213-24-5
Polyvinyl alcohol	9002-89-5
Polyvinyl chloride	None
Polyvinyl chloride, minimum number average molecular weight (in amu), 29,000	9002-86-2
Poly(vinylpyrrolidone), minimum number average molecular weight (in amu), 4,000	9003-39-8
Poly(vinylpyrrolidone-1-eicosene), minimum average molecular weight (in amu), 3,000	28211-18-9
Poly(vinylpyrrolidone-1-hexadecene), minimum average molecular weight (in amu), 4,700	63231-81-2
1-Propanesulfonic acid, 2-methyl-2-[(1-oxo-2-propenyl)amino]-, monosodium salt, polymer with ethenol and ethenyl acetate, minimum number average molecular weight (in amu) 50,000	107568-12-7
2-Propene-1-sulfonic acid sodium salt, polymer with ethenol and ethenyl acetate, number average molecular weight (in amu) 6,000-12,000	None
2-Propenoic acid, butyl ester, polymer with ethyl 2-propenoate and N-(hydroxymethyl)-2-propenamamide, minimum number average molecular weight (in amu), 30,000	33438-19-6
2-Propenoic acid, methyl ester, polymer with ethenyl acetate, hydrolyzed, sodium salts.	886993-11-9
2-Propenoic acid, 2-Methyl-, Polymer with Butyl 2-Propenoate, Methyl 2-Methyl-2-Propenoate, Methyl 2-Propenoate and 2-Propenoic Acid, graft, Compound with 2-Amino-2-Methyl-1-Propanol	153163-36-1
2-Propenoic Acid, 2-Methyl-, Polymer with Ethenylbenzene, 2-Ethylhexyl 2-Propenoate, 2-Hydroxyethyl 2-Propenoate, N-(Hydroxymethyl) 2-Methyl-2-Propenamamide and Methyl 2-Methyl-2-Propenoate, Ammonium Salt	146753-99-3

Environmental Protection Agency

§ 180.960

Polymer	CAS No.
2-Propenoic, 2-methyl-, polymers with ethyl acrylate and polyethylene glycol methylacrylate C ₁₈₋₂₂ alkyl ethers	888969-14-0
2-Propenoic acid, 2-hydroxyethyl ester, polymer with α -[4-(ethenyloxy)butyl]- ω -hydroxypoly (oxy-1,2-ethanediyl), minimum number average molecular weight (in amu), 17,000	1007234-89-0
2-Propenoic acid, monoester with 1,2-propanediol, polymer with α -[4-(ethenyloxy) butyl]- ω -hydroxypoly (oxy-1,2-ethanediyl) and 2,5-furandione, minimum number average molecular weight (in amu), 25,000	955015-23-3
2-Propenoic acid, polymer with α -[4-(ethenyloxy) butyl]- ω -hydroxypoly (oxy-1,2-ethanediyl) and 2,5-furandione, sodium salt, minimum number average molecular weight (in amu), 25,000	251479-97-7
2-Propenoic acid, polymer with α -[4-(ethenyloxy) butyl]- ω -hydroxypoly (oxy-1,2-ethanediyl) and 1,2-propanediol mono-2-propenoate, potassium sodium salt, minimum number average molecular weight (in amu), 16,000	518026-64-7
2-Propenoic acid, polymer with α -[4-(ethenyloxy) butyl]- ω -hydroxypoly (oxy-1, 2-ethanediyl), sodium salt, minimum number average molecular weight (in amu), 24,000	250591-84-5
2-Propenoic acid, polymer with 2-propenamide, sodium salt, minimum number average molecular weight (in amu), 18,000	25085-02-3
2-Propenoic acid, sodium salt, polymer with 2-propenamide, minimum number average molecular weight (in amu), 18,000	25987-30-8
Silane, dichloromethyl- reaction product with silica minimum number average molecular weight (in amu), 3,340,000	68611-44-9
Silane, trimethoxy[3-(oxiranylmethoxy)propyl]-, hydrolysis products with silica, minimum number average molecular weight (in amu), 640,000	68584-82-7
Sodium polyflavonoidsulfonate, consisting chiefly of the copolymer of catechin and leucocyanidin	None
Soybean oil, ethoxylated; the poly(oxyethylene) content averages 10 moles or greater	61791-23-9
Starch, oxidized, polymers with Bu acrylate, tert-Bu acrylate and styrene, minimum number average molecular weight (in amu), 10,000	204142-80-3
Stearyl methacrylate-1,6-hexanediol dimethacrylate copolymer, minimum molecular weight (in amu), 100,000	None

Polymer	CAS No.
Styrene, copolymers with acrylic acid and/or methacrylic acid, with none and/or one or more of the following monomers: Acrylamidopropyl methyl sulfonic acid, methallyl sulfonic acid, 3-sulfopropyl acrylate, 3-sulfopropyl methacrylate, hydroxypropyl methacrylate, hydroxypropyl acrylate, hydroxyethyl methacrylate, and/or hydroxyethyl acrylate; and its sodium, potassium, ammonium, monoethanolamine, and triethanolamine salts; the resulting polymer having a minimum number average molecular weight (in amu), 1,200	None
Styrene, 2-ethylhexyl acrylate, butyl acrylate copolymer, minimum number average molecular weight (in amu), 4,200	30795-23-4
Styrene-2-ethylhexyl acrylate-glycidyl methacrylate-2-acrylamido-2-methylpropanesulfonic acid graft copolymer, minimum number average molecular weight (in amu), 12,500	None
Styrene-maleic anhydride copolymer	None
Styrene-maleic anhydride copolymer, ester derivative	None
Tetradecyl acrylate-acrylic acid copolymer, minimum number average molecular weight (in amu), 3,000	None
Tetraethoxysilane, polymer with hexamethyldisiloxane, minimum number average molecular weight (in amu), 6,500	104133-09-7
α -[p-(1,1,3,3-Tetramethylbutyl)phenyl]- ω -hydroxypoly(oxyethylene) produced by the condensation of 1 mole of p-(1,1,3,3-tetramethylbutyl)phenol with a range of 30-70 moles of ethylene oxide	9036-19-5 9002-93-1
α -[p-(1,1,3,3-Tetramethylbutyl)phenyl] poly(oxypropylene) block polymer with poly(oxyethylene); the poly(oxypropylene) content averages 25 moles, the poly(oxyethylene) content averages 40 moles, the molecular weight (in amu) averages 3,400	None
α -[2,4,6-Tris[1-(phenyl)ethyl]phenyl]- ω -hydroxy poly(oxyethylene) poly(oxypropylene) copolymer, the poly(oxypropylene) content averages 2-8 moles, the poly(oxyethylene) content averages 16-30 moles, average molecular weight (in amu), 1,500	None
Urea-formaldehyde copolymer, minimum average molecular weight (in amu), 30,000	9011-05-6
Vinyl acetate-allyl acetate-monomethyl maleate copolymer, minimum average molecular weight (in amu), 20,000	None
Vinyl acetate-ethylene copolymer, minimum number average molecular weight (in amu), 69,000	24937-78-8

§ 180.1011

40 CFR Ch. I (7–1–09 Edition)

Polymer	CAS No.
Vinyl acetate polymer with none and/or one or more of the following monomers: Ethylene, propylene, N-methyl acrylamide, acrylamide, monoethyl maleate, diethyl maleate, monoethyl maleate, dioctyl maleate, maleic anhydride, maleic acid, octyl acrylate, butyl acrylate, ethyl acrylate, methyl acrylate, acrylic acid, octyl methacrylate, butyl methacrylate, ethyl methacrylate, methyl methacrylate, methacrylic acid, carboxyethyl acrylate, and diallyl phthalate; and their corresponding sodium, potassium, ammonium, isopropylamine, triethylamine, monoethanolamine and/or triethanolamine salts; the resulting polymer having a minimum number average molecular weight (in amu), 1,200	None
Vinyl acetate-vinyl alcohol-alkyl lactone copolymer, minimum number average molecular weight (in amu), 40,000; minimum viscosity of 18 centipoise	None
Vinyl alcohol-disodium itaconate copolymer, minimum average molecular weight (in amu), 50,290	None
Vinyl alcohol-vinyl acetate copolymer, benzaldehyde-o-sodium sulfonate condensate, minimum number average molecular weight (in amu), 20,000	None
Vinyl alcohol-vinyl acetate-monomethyl maleate, sodium salt-maleic acid, disodium salt-γ-butyrolactone acetic acid, sodium salt copolymer, minimum number average molecular weight (in amu), 20,000	None
Vinyl chloride-vinyl acetate copolymers	None
Vinyl pyrrolidone-acrylic acid copolymer, minimum number average molecular weight (in amu), 6,000	28062–44–4
Vinyl pyrrolidone-dimethylaminoethylmethacrylate copolymer, minimum number average molecular weight (in amu), 20,000	30581–59–0
Vinyl pyrrolidone-styrene copolymer	25086–29–7

[67 FR 36528, May 24, 2002]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 180.960, see the List of CFR Sections Affected, which appears in the Finding Aids section of the printed volume and on GPO Access.

§ 180.1011 Viable spores of the microorganism *Bacillus thuringiensis* Berliner; exemption from the requirement of a tolerance.

(a) For the purposes of this section the microbial insecticide for which exemption from the requirement of a tolerance is being established shall have the following specifications:

(1) The microorganism shall be an authentic strain of *Bacillus thuringiensis* Berliner conforming to the morphological and biochemical characteristics of *Bacillus thuringiensis* as described in Bergey's Manual of Determinative Bacteriology, Eighth Edition.

(2) Spore preparations of *Bacillus thuringiensis* Berliner shall be produced by pure culture fermentation procedures with adequate control measures during production to detect any changes from the characteristics of the parent strain or contamination by other microorganisms.

(3) Each lot of spore preparation, prior to the addition of other materials, shall be tested by subcutaneous injection of at least 1 million spores into each of five laboratory test mice weighing 17 grams to 23 grams. Such test shall show no evidence of infection or injury in the test animals when observed for 7 days following injection.

(4) Spore preparations shall be free of the *Bacillus thuringiensis* β-exotoxin when tested with the fly larvae toxicity test ("Microbial Control of Insects and Mites," R.P.M. Bond et al., p. 280 ff., 1971). This specification can be satisfied either by determining that each master seed lot brought into production is a *Bacillus thuringiensis* strain which does not produce β-exotoxin under standard manufacturing conditions or by periodically determining that β-exotoxin synthesized during spore production is eliminated by the subsequent spore-harvesting procedure.

(b) Exemption from the requirement of a tolerance is established for residues of the microbial insecticide *Bacillus thuringiensis* Berliner, as specified in paragraph (a) of this section, in or on honey and honeycomb and all other raw agricultural commodities when it is applied either to growing crops, or when it is applied after harvest in accordance with good agricultural practices.

[36 FR 22540, Nov. 25, 1971, as amended at 38 FR 19045, July 17, 1973; 42 FR 28540, June 3, 1977; 45 FR 43721, June 30, 1980; 45 FR 56347, Aug. 25, 1980; 74 FR 26533, June 3, 2009]