

§ 558.367 Niclosamide.

(a) *Approvals.* Type A medicated articles: 66 percent to 000859 in § 510.600(c) of this chapter.

(b) *Conditions of use.* It is used in the feed of laboratory mice as follows:

(1) *Amount per ton.* 6.6 pounds (0.33 percent).

(2) *Indications for use.* Control of the mouse tapeworm *Hymenolepis nana*.

(3) *Limitations.* Feed daily for 7 consecutive days. Do not administer other drugs during treatment. If reinfection occurs, treatment may be repeated after 30 days. Do not feed to sick or debilitated animals or animals under severe stress. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism. When mixed according to directions and fed free choice, the feed will provide a daily dose of 500 milligrams per kilogram of body weight.

[41 FR 36810, Sept. 1, 1976, as amended at 51 FR 7399, Mar. 3, 1986]

§ 558.369 Nitarsone.

(a) *Approvals.* Type A medicated articles: 50 percent to 046573 in § 510.600(c) of this chapter.

(b) *Related tolerances.* See § 556.60 of this chapter.

(c) *NAS/NRC status.* These conditions of use are NAS/NRC reviewed and found effective. NADA's for these uses may not require effectiveness data as specified by § 514.111 of this chapter, but may require bioequivalency and safety information.

(d) *Conditions of use.* It is used for chickens and turkeys as follows:

(1) *Amount.* Nitarsone, 0.01875 percent.

(2) *Indications for use.* As an aid in the prevention of blackhead.

(3) *Limitations.* Early medication is essential to prevent spread of disease. Adequate drinking water must be provided near feeder at all times. The drug is not effective in preventing blackhead in birds infected more than 4 or 5 days. Discontinue use 5 days before slaughtering animals for human consumption to allow elimination of the drug from edible tissues. The drug is dangerous for ducks, geese, and dogs. Overdosage or lack of water may result

in leg weakness or paralysis. Use as sole source of arsenic.

[46 FR 47535, Sept. 29, 1981, as amended at 47 FR 14152, Apr. 2, 1982; 51 FR 7399, Mar. 3, 1986; 52 FR 2686, Jan. 26, 1987; 55 FR 8460, Mar. 8, 1990; 57 FR 8578, Mar. 11, 1992]

§ 558.376 Nitromide and sulfanitran.

(a) *Approvals.* Type A medicated articles: 25 percent nitromide, 30 percent sulfanitran, with or without 5 percent roxarsone to 053501 in § 510.600(c) of this chapter.

(b) *Related tolerances.* See §§ 556.220 and 556.680 of this chapter.

(c) *Conditions of use.* It is used for chickens as follows:

(1) *Amount.* 227 grams per ton nitromide (0.025 percent) and 272 grams per ton sulfanitran (0.03 percent).

(i) *Indications for use.* As an aid in the prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, and *E. acervulina*.

(ii) *Limitations.* Not to be fed to laying chickens; withdraw 5 days before slaughter; from Type A articles containing not more than 25 percent nitromide and 30 percent sulfanitran.

(2) *Amount.* 227 grams per ton nitromide (0.025 percent) and 272 grams per ton sulfanitran (0.03 percent), plus 45.4 grams per ton roxarsone (0.005 percent).

(i) *Indications for use.* Prevention of coccidiosis caused by *Eimeria tenella*, *E. necatrix*, and *E. acervulina*; growth promotion and feed efficiency; improving pigmentation.

(ii) *Limitations.* Not to be fed to laying chickens; withdraw 5 days before slaughter; from Type A articles containing not more than 25 percent nitromide, 30 percent sulfanitran, and 5 percent roxarsone; as sole source of organic arsenic.

[41 FR 11002, Mar. 15, 1976, as amended at 51 FR 7399, Mar. 3, 1986; 52 FR 2686, Jan. 26, 1987; 55 FR 8460, Mar. 8, 1990]

§ 558.415 Novobiocin.

(a) *Approvals.* Type A medicated articles: 25 grams of activity per pound to 000009 in § 510.600(c) of this chapter. Type B medicated feeds: 17.5 grams per pound to 000009 in § 510.600(c) of this chapter.

(b) *Related tolerances.* See § 556.460 of this chapter.