

**Food and Drug Administration, HHS**

**§ 369.21**

*Caution*—Do not use in the eyes or nose. Not for prolonged use. Do not apply to large areas of the body. If redness, irritation, swelling, or pain persists or increases, discontinue use unless directed by a physician.

SODIUM GENTISATE. (See §§ 201.314 and 310.301(a)(2) of this chapter.)

*Warning*—Do not use in children under 6 years of age or use for prolonged period unless directed by physician.

“Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.”

If offered for use in arthritis or rheumatism, in juxtaposition therewith, the statement:

*Caution*—If pain persists for more than 10 days, or redness is present, or in conditions affecting children under 12 years of age, consult a physician immediately.

TUAMINOHEPTANE SULFATE NASAL PREPARATIONS. (See § 310.201(a)(16) of this chapter.)

*Caution*—Do not exceed recommended dosage. Overdosage may cause nervous-

ness, restlessness, or sleeplessness. Individuals with high blood pressure, heart disease, diabetes, or thyroid disease should use only as directed by physician. Do not use for more than 3 or 4 consecutive days unless directed by physician.

VIBESATE PREPARATIONS. (See § 310.201(a)(18) of this chapter.)

*Caution*—Do not use but consult physician for deep or puncture wounds or serious burns. If redness, irritation, swelling, or pain persists or increases, discontinue use and consult physician.

*Warning*—Contents under pressure. Do not puncture. Do not use or store near heat or open flame. Exposure to temperatures above 130° Fahrenheit may cause bursting. Never throw container into fire or incinerator.

[39 FR 11745, Mar. 29, 1974]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 369.21, see the List of CFR Sections Affected, which appears in the Finding Aids section of the printed volume and on GPO Access.

**PARTS 370–499 [RESERVED]**