## Food and Drug Administration, HHS

established in veal calves. A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for yeal.

- (3) Steers fed in confinement for slaughter—(i) Amount. Reimplant 200 mg progesterone and 20 mg estradiol benzoate on approximately day 70 following an initial implant of 100 mg progesterone and 10 mg estradiol benzoate or 200 mg progesterone and 20 mg estradiol benzoate.
- (ii) *Indications for use*. For additional improvement in rate of weight gain.
- (iii) Limitations. For subcutaneous ear implantation. Safety and effectiveness have not been established in veal calves. A withdrawal period has not been established for this product in preruminating calves. Do not use in calves to be processed for veal.

[69 FR 70055, Dec. 2, 2004]

## § 522.1962 Promazine hydrochloride.

- (a) *Specifications*. Each milliliter of solution contains 50 milligrams (mg) promazine hydrochloride.
- (b) *Sponsors*. See sponsors in §510.600(c) of this chapter for use as in paragraph (c) of this section:
- (1) No. 000856 for use as in paragraphs (c)(1)(i)(A), (c)(1)(ii)(A), (c)(1)(iii), and (c)(2) of this section.
- (2) No. 061623 for use as in paragraphs (c)(1)(i)(B), (c)(1)(ii)(B), and (c)(1)(iii) of this section.
- (c) Conditions of use—(1) Horses—(i) Amount—(A) 0.2 to 0.5 milligrams per pounds (mg/lb) body weight intramuscularly or intravenously every 4 to 6 hours.
- (B) 0.2 to 0.5 mg/lb body weight intravenously as required.
- (ii) Indications for use—(A) For use as a tranquilizer, preanesthetic, or for minor operative procedures in conjunction with local anesthesia; and as adjunctive therapy for tetanus.
- (B) For use as a tranquilizer and preanesthetic.
- (iii) *Limitations*. Not for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.
- (2) Dogs and cats—(i) Amount. 1 to 2 mg/lb body weight intramuscularly or intravenously every 4 to 6 hours.

- (ii) Indications for use. For use as a tranquilizer, preanesthetic, for minor operative procedures in conjunction with local anesthesia, as adjunctive therapy for tetanus, and as an antiemetic prior to worming; or to prevent motion sickness in dogs.
- (iii) *Limitations*. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[46 FR 18962, Mar. 27, 1981, as amended at 68 FR 59881, Oct. 20, 2003; 70 FR 50183, Aug. 26, 2005]

## § 522.2002 Propiopromazine hydrochloride injection.

- (a) Chemical name. 1-Propanone, 1-[10-[3-(dimethylamino) propyl] phenothiazine-2-yl]-, monohydrochloride.
- (b) Specifications. Propiopromazine hydrochloride injection contains 5 or 10 milligrams of the drug in each milliliter of sterile aqueous solution.
- (c) *Sponsor*. See No. 000856 in §510.600(c) of this chapter.
- (d) Conditions of use. (1) It is administered either intravenously or intramuscularly to dogs and cats for tranquilization at a dosage level of 0.05–0.5 milligram per pound of body weight and is also administered intravenously to dogs and cats as a preanesthetic at a dosage level of 0.25 milligram per pound of body weight.
- (2) It is not to be used in conjunction with organophosphates and/or procaine hydrochloride since phenothiazines may potentiate the toxicity of organophosphates and the activity of procaine hydrochloride.
- (3) For use only by or on the order of a licensed veterinarian.

[40 FR 13858, Mar. 27, 1975, as amended at 46 FR 60570, Dec. 11, 1981; 61 FR 5507, Feb. 13, 1996]

## §522.2005 Propofol injection.

- (a) Specifications. The drug is a sterile, nonpyrogenic, oil-in-water emulsion containing 10 milligrams of propofol per milliliter.
- (b) Sponsor. See No. 000061 in  $\S510.600(c)$  of this chapter for use as in paragraphs (c)(1) and (c)(2) of this section. See No. 000074 in  $\S510.600(c)$  of this chapter for use as in paragraph (c)(1) of this section.