§ 529.1044

dissolved oxygen is less than 5 milligrams per liter.

[51 FR 11441, Apr. 3, 1986, as amended at 58 FR 59169, Nov. 8, 1993; 59 FR 60076, Nov. 22, 1994; 63 FR 38304, July 16, 1998; 68 FR 5563; Feb. 4, 2003; 72 FR 45158, Aug. 13, 2007]

§ 529.1044 Gentamicin sulfate in certain other dosage forms.

§ 529.1044a Gentamicin sulfate intrauterine solution.

- (a) Specifications. Each milliliter of solution contains 50 or 100 milligrams gentamicin sulfate.
- (b) *Sponsors*. See Nos. 000010, 000061, 000856, 057561, 058005, 059130, and 061623 in §510.600(c) of this chapter.
- (c) Conditions of use in horses—(1) Amount. Infuse 2 to 2.5 grams per day for 3 to 5 days during estrus.
- (2) Indications for use. For control of bacterial infections of the uterus (metritis) and as an aid in improving conception in mares with uterine infections caused by bacteria sensitive to gentamicin.
- (3) *Limitations*. Do not use in horses intended for human consumption. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

§ 529.1044b Gentamicin sulfate solution.

[71 FR 51727, Aug. 31, 2006]

- (a) Specifications. Each milliliter of solution contains gentamicin sulfate equivalent to 50 milligrams of gentamicin base.
- (b) *Sponsors*. See Nos. 000061 and 054925 in §510.600(c) of this chapter.
- (c) Conditions of use. (1) The drug is recommended as an aid in the reduction or elimination of the following microorganisms from turkey-hatching eggs: Arizona hinshawii (paracolon), Salmonella st. paul, and Mycoplasma meleagridis.
- (2) The drug is added to clean water to provide a dip solution with a gentamicin concentration of 250 to 1,000 parts per million. A concentration of 500 parts per million is recommended. Clean eggs should be held submerged in the gentamicin solution under a vacuum of about 27.5 to 38 centimeters of mercury for 5 minutes followed by additional soaking in gentamicin solution for approximately

10 minutes at atmospheric pressure. Eggs can also be treated by warming them for 3 to 6 hours at approximately 100 °F. then immediately submerging them in gentamicin solution maintained at about 40 °F., keeping the eggs submerged for 10 to 15 minutes.

(3) For use in the dipping treatment of turkey-hatching eggs only. Eggs which have been dipped in the drug shall not be used for food.

[40 FR 13881, Mar. 27, 1975, as amended at 52 FR 7833, Mar. 13, 1987; 62 FR 22889, Apr. 28, 1997; 71 FR 13543, Mar. 16, 2006]

§ 529.1115 Halothane.

- (a) *Specifications*. The drug is a colorless, odorless, nonflammable, nonexplosive, heavy liquid containing 0.01 percent thymol as a preservative.
- (b) Sponsor. See 000856 and 012164 in \$510.600(c) of this chapter.
- (c) Conditions of use—(1) Amount. Two to 5 percent of inhaled atmosphere for induction of anesthesia; 0.5 to 2 percent for maintenance of anesthesia.¹
- (2) Indications for use. For nonfood animals for the induction and maintenance of anesthesia.¹
- (3) Limitations. Administered by inhalation. May be administered with either oxygen or a mixture of oxygen and nitrous oxide. Place drug vaporizer between the gas supply and breathing bag to prevent overdosage. Not recommended for obstetrical anesthesia except when uterine relaxation is required. Do not use in pregnant animals; information on possible adverse effects on fetal development is not available. Operating rooms should have adequate ventilation to prevent accumulation of anesthetic gases. Not for use in animals intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.1

[46 FR 27915, May 22, 1981, as amended at 62 FR 29014, May 29, 1997]

§529.1150 Hydrogen peroxide.

(a) Specifications. Each milliliter of solution contains 396.1 milligrams (mg)

¹These conditions have been reviewed by FDA and found effective. NADA's for similar products for these conditions of use need not include effectiveness data as specified by §514.111 of this chapter, but may require bioequivalency and safety information.