

## § 556.520

### § 556.520 Prednisolone.

A tolerance of zero is established for residues of prednisolone in milk from dairy animals.

### § 556.530 Prednisone.

A tolerance of zero is established for residues of prednisone in milk from dairy animals.

### § 556.540 Progesterone.

No residues of progesterone are permitted in excess of the following increments above the concentrations of progesterone naturally present in untreated animals:

(a) In uncooked edible tissues of steers and calves:

- (1) 3 parts per billion for muscle.
- (2) 12 parts per billion for fat.
- (3) 9 parts per billion for kidney.
- (4) 6 parts per billion for liver.

(b) In uncooked edible tissues of lambs:

- (1) 3 parts per billion for muscle.
- (2) 15 parts per billion for fat, kidney, and liver.

[49 FR 13873, Apr. 9, 1984]

### § 556.550 Propylparaben.

A tolerance of zero is established for residues of propylparaben in milk from dairy animals.

### § 556.560 Pyrantel tartrate.

Tolerances are established for residues of pyrantel tartrate in edible tissues of swine as follows:

- (a) 10 parts per million in liver and kidney.
- (b) 1 part per million in muscle.

### § 556.570 Ractopamine.

(a) *Acceptable Daily Intake (ADI)*. The ADI for total residues of ractopamine hydrochloride is 1.25 micrograms per kilogram of body weight per day.

(b) *Tolerances*—(1) *Cattle*—(i) *Liver (the target tissue)*. The tolerance for ractopamine hydrochloride (the marker residue) is 0.09 parts per million (ppm).

(ii) *Muscle*. The tolerance for ractopamine hydrochloride (the marker residue) is 0.03 ppm.

(2) *Swine*—(i) *Liver (the target tissue)*. The tolerance for ractopamine hydro-

## 21 CFR Ch. I (4–1–10 Edition)

chloride (the marker residue) is 0.15 ppm.

(ii) *Muscle*. The tolerance for ractopamine hydrochloride (the marker residue) is 0.05 ppm.

(3) *Turkeys*—(i) *Liver (the target tissue)*. The tolerance for ractopamine (the marker residue) is 0.45 ppm.

(ii) *Muscle*. The tolerance for ractopamine (the marker residue) is 0.1 ppm.

[68 FR 54659, Sept. 18, 2003, as amended at 73 FR 72715, Dec. 1, 2008]

### § 556.580 Robenidine hydrochloride.

Tolerances are established for residues of robenidine hydrochloride in edible tissues of chickens as follows:

(a) 0.2 part per million in skin and fat.

(b) 0.1 part per million (negligible residue) in edible tissues other than skin and fat.

### § 556.590 Salicylic acid.

A tolerance of zero is established for residues of salicylic acid in milk from dairy animals.

### § 556.592 Salinomycin.

(a) *Acceptable daily intake (ADI)*. The ADI for total residues of salinomycin is 0.005 milligram per kilogram of body weight per day.

(b) [Reserved]

[65 FR 70791, Nov. 28, 2000]

### § 556.597 Semduramicin.

(a) *Acceptable daily intake (ADI)*. The ADI for total residues of semduramicin is 180 micrograms per kilogram of body weight per day.

(b) *Tolerances*—(1) *Broiler chickens*. Tolerances are established for residues of parent semduramicin in uncooked edible tissues of 400 parts per billion (ppb) in liver and 130 ppb in muscle.

(2) [Reserved]

[64 FR 48296, Sept. 3, 1999]

### § 556.600 Spectinomycin.

(a) *Acceptable daily intake (ADI)*. The ADI for total residues of spectinomycin is 25 micrograms per kilogram of body weight per day.

(b) *Chickens and turkeys*. A tolerance of 0.1 part per million (ppm) for negligible residues of spectinomycin in