Food and Drug Administration, HHS

Pt. 558

§556.745 Tulathromycin.

(a) Acceptable daily intake (ADI). The ADI for total residues of tulathromycin is 15 micrograms per kilogram of body weight per day.

(b) *Tolerances*—(1) *Cattle*—(i) *Liver* (*the target tissue*). The tolerance for CP– 60,300 (the marker residue) is 5.5 parts per million (ppm).

(ii) [Reserved]

(2) Swine—(i) Kidney (the target tissue). The tolerance for CP-60,300 (the marker residue) is 15 ppm.

(ii) [Reserved]

(c) *Related conditions of use*. See §522.2630 of this chapter.

[70 FR 39918, July 12, 2005]

§556.750 Virginiamycin.

(a) Acceptable daily intake (ADI). The ADI for total residues of virginiamycin is 250 micrograms per kilogram of body weight per day.

(b) *Tolerances*—(1) *Swine*. Tolerances are established for residues of virginiamycin in uncooked edible tissues of 0.4 part per million (ppm) in kidney, skin, and fat, 0.3 ppm in liver, and 0.1 ppm in muscle.

(2) Broiler chickens and cattle. A tolerance for residues of virginiamycin is not required.

[64 FR 48296, Sept. 3, 1999]

§ 556.760 Zeranol.

(a) Acceptable daily intake (ADI). The ADI for total residues of zeranol is 0.00125 milligrams per kilogram of body weight per day.

(b) *Tolerances*. The tolerances for residues of zeranol in edible tissues are:

(1) Cattle. A tolerance is not needed.

(2) Sheep. 20 parts per billion.

(c) *Related conditions of use*. See §522.2680 of this chapter.

[40 FR 13942, Mar. 27, 1975, as amended at 54
FR 31950, Aug. 3, 1989; 67 FR 6867, Feb. 14, 2002; 70 FR 15759, Mar. 29, 2005]

§556.765 Zilpaterol.

(a) Acceptable daily intake (ADI). The ADI for total residues of zilpaterol is 0.083 micrograms per kilogram of body weight per day.

(b) Tolerances—(1) Cattle—(i) Liver (the target tissue). The tolerance for zilpaterol freebase (the marker residue) is 12 parts per billion (ppb).

(ii) [Reserved]

(2) [Reserved]

[71 FR 53005, Sept. 8, 2006]

§556.770 Zoalene.

Tolerances are established for residues of zoalene (3,5-dinitro-*o*toluamide) and its metabolite 3-amino-5-nitro-*o*-toluamide in food as follows:

(a) In edible tissues of chickens:

(1) 6 parts per million in uncooked liver and kidney.

(2) 3 parts per million in uncooked muscle tissue.

(3) 2 parts per million in uncooked fat.

(b) In edible tissues of turkeys: 3 parts per million in uncooked muscle tissue and liver.

PART 558—NEW ANIMAL DRUGS FOR USE IN ANIMAL FEEDS

Subpart A—General Provisions

Sec.

- 558.3 Definitions and general considerations applicable to this part.
- 558.4 Requirement of a medicated feed mill license.
- 558.5 Requirements for liquid medicated feed.
- 558.6 Veterinary feed directive drugs.
- 558.15 Antibiotic, nitrofuran, and sulfonamide drugs in the feed of animals.

Subpart B—Specific New Animal Drugs For Use in Animal Feeds

- 558.35 Aklomide.
- 558.55 Amprolium.
- 558.58 Amprolium and ethopabate.
- 558.59 Apramycin.
- 558.60 Arsanilate sodium.
- 558.62 Arsanilic acid.
- 558.76 Bacitracin methylene disalicylate.
- 558.78 Bacitracin zinc. 558.95 Bambermycins
- 558.95 Bambermycins. 558.105 [Reserved]
- 558 115 Carbadox
- 558.120 Carbarsone (not U.S.P.).
- 558.128 Chlortetracycline.
- 558.140 Chlortetracycline
- sulfamethazine. 558.145 Chlortetracycline, procaine penicillin, and sulfamethazine.

and

- 558.155 Chlortetracycline, sulfathiazole, penicillin.
- 558.175 Clopidol.
- 558.185 Coumaphos.
- 558.195 Decoquinate.