Environmental Protection Agency

field, hay; ryegrass, Italian, hay; sorghum, grain, grain; soybean, hay; sudangrass, forage; sudangrass, hay; timothy, forage; timothy, hay; vetch, forage; vetch, hay; and wheat, grain.

(b) Propanoic acid is exempt from the requirement of a tolerance for residues in or on cattle, meat; cattle, meat byproducts; goat, meat; goat, meat byproducts; horse, meat; horse, meat byproducts; horse, meat; horse, meat byproducts; sheep, meat; sheep meat byproducts; and, poultry, fat; poultry meat; poultry meat byproducts; and, and egg when applied as a bactericide/fungicide to livestock drinking water, poultry litter, and storage areas for silage and grain.

(c) Preharvest and postharvest application of propanoic acid (CAS Reg. No. 79–09–4), propanioc acid, calcium salt (CAS Reg. No. 4075–81–4), and propanioc sodium salt (CAS Reg. No. 137–40–6) are exempted from the requirement of a tolerance on all crops when used as either an active or inert ingredient in accordance with good agricultural practice in pesticide formulations applied to growing crops, to raw agricultural commodities before and after harvest and to animals.

[69 FR 47025, Aug. 4, 2004, as amended at 74 FR 26534, June 3, 2009]

§180.1025 Xylene; exemption from the requirement of a tolerance.

Xylene is exempted from the requirement of a tolerance when used as an aquatic herbicide applied to irrigation conveyance systems in accordance with the following conditions:

- (a) It is to be used only in programs of the Bureau of Reclamation, U.S. Department of Interior, and cooperating water user organizations.
- (b) It is to be applied as an emulsion at an initial concentration not to exceed 750 parts per million.
- (c) It is not to be applied when there is any likelihood that the irrigation water will be used as a source of raw water for a potable water system or where return flows of such treated irrigation water into receiving rivers and streams would contain residues of xylene in excess of 10 parts per million.
- (d) Xylene to be used as an aquatic herbicide shall meet the requirement limiting the presence of a polynuclear

aromatic hydrocarbons as listed in 21 CFR 172.250.

[38 FR 16352, June 22, 1973, as amended at 50 FR 2980, Jan. 3, 1985]

§180.1027 Nuclear polyhedrosis virus of Heliothis zea; exemption from the requirement of a tolerance.

- (a) For the purposes of this section, the viral insecticide must be produced with an unaltered and unadulterated inoculum of the single-embedded *Heliothis zea* nuclear polyhedrosis virus (HzSNPV). The identity of the seed virus must be assured by periodic checks.
- (b) Each lot of active ingredient of the viral insecticide shall have the following specifications:
- (1) The level of extraneous bacterial contamination of the final unformulated viral insecticide should not exceed 10⁷ colonies per gram as determined by an aerobic plate on trypticase soy agar.
- (2) Human pathogens, e.g., Salmonella, Shigella, or Vibrio, must be absent.
- (3) Safety to mice as determined by an intraperitoneal injection study must be demonstrated.
- (4) Identity of the viral product, as determined by the most sensitive and standardized analytical technique, e.g., restriction endonuclease and/or SDS-PAGE analysis, must be demonstrated.
- (c) Exemptions from the requirement of a tolerance are established for the residues of the microbial insecticide *Heliothis zea* NPV, as specified in paragraphs (a) and (b) of this section, in or on all agricultural commodities.

[60 FR 42460, Aug. 16, 1995, as amended at 74 FR 26534, June 3, 2009]

§ 180.1033 Methoprene; exemption from the requirement of a tolerance.

Methoprene is exempt from the requirement of a tolerance in or on all food commodities when used to control insect larvae.

[68 FR 34829, June 11, 2003]

§ 180.1035 Pine oil; exemption from the requirement of a tolerance.

Pine oil is exempted from the requirement of a tolerance for residues in