

§ 180.473

40 CFR Ch. I (7-1-10 Edition)

Commodity	Parts per million
Raspberry, wild	2.5
Safflower, seed	0.05
Salal	3.5
Sapodilla	1.0
Sapote, black	1.0
Sapote, mamey	1.0
Sheep, fat	0.30
Sheep, meat	0.30
Sheep, meat byproducts	0.30
Soursop	0.30
Soybean, forage	8.0
Soybean, hay	35
Soybean, meal	4.0
Soybean, seed	3.5
Spanish lime	3.0
Star apple	1.0
Starfruit	1.0
Strawberry	0.50
Sugar apple	0.30
Sunflower, seed	0.05
Tomato, paste	6.0
Tomato, puree	3.0
Vegetable, brassica leafy, group 5	3.5
Vegetable, cucurbit, group 9	0.5
Vegetable, fruiting, group 8	1.0
Vegetable, leaves of root and tuber, group 2	4.0
Vegetable, legume, group 6, except soybean	4.0
Vegetable, root and tuber, group 1, except sugar beet	0.40
Watercress	3.5
Watercress, upland	3.5
Wax jambu	1.0

(b) Section 18 emergency exemptions. [Reserved]

(c) Tolerances with regional registrations. [Reserved]

(d) Indirect or inadvertent residues. Tolerances are established for indirect or inadvertent residues of the insecticide imidacloprid, including its metabolites and degradates, in or on the commodities in the table below. Compliance with the tolerance levels specified below is to be determined by measuring only the sum of imidacloprid (1-[6-chloro-3-pyridinyl] methyl)-N-nitro-2-imidazolidinimine) and its metabolites containing the 6-chloropyridinyl moiety, calculated as the stoichiometric equivalent of imidacloprid, in or on the following commodities, when present therein as a result of the application of the pesticide to growing crops listed in this section and other non-food crops as follows:

Commodity	Parts per million
Rice, grain	0.05
Vegetable, foliage of legume, group 7	2.5
Vegetable, legume, group 6	0.3

§ 180.473 Glufosinate ammonium; tolerances for residues.

(a) General. Tolerances are established for residues of the herbicide glufosinate-ammonium (butanoic acid, 2-amino-4-(hydroxymethylphosphinyl)monoammonium salt) and its metabolites, 2-acetamido-4-methylphosphinico-butanoic acid and 3-methylphosphinico-propionic acid, expressed as 2-amino-4-(hydroxymethylphosphinyl)butanoic acid equivalents, in or on the following food commodities:

Commodity	Parts per million
Almond, hulls	0.50
Apple	0.05
Banana	0.30
Banana, pulp	0.20
Beet, sugar, molasses	5.0
Beet, sugar, roots	0.9
Beet, sugar, tops (leaves)	1.5
Bushberry subgroup 13B	0.15
Canola, meal	1.1
Canola, seed	0.40
Cattle, fat	0.40
Cattle, meat	0.15
Cattle, meat byproducts	6.0
Corn, field forage	4.0
Corn, field, grain	0.20
Corn, field, stover	6.0
Cotton, gin byproducts	15
Cotton, undelinted seed	4.0
Egg	0.15
Goat, fat	0.40
Goat, meat	0.15
Goat, meat byproducts	6.0
Grain aspirated fractions	25
Grape	0.05
Hog, fat	0.40
Hog, meat	0.15
Hog, meat byproducts	6.0
Horse, fat	0.40
Horse, meat	0.15
Horse, meat byproducts	6.0
Juneberry	0.10
Lingonberry	0.10
Milk	0.15
Nut, tree, group 14	0.10
Pistachio	0.10
Potato	0.80
Potato, chips	1.6
Potato granules/flakes	2.0
Poultry, fat	0.15
Poultry, meat	0.15
Poultry, meat byproducts	0.60
Rice, grain	1.0
Rice, hull	2.0
Rice, straw	2.0
Salal	0.10
Sheep, fat	0.40
Sheep, meat	0.15
Sheep, meat byproducts	6.0
Soybean	2.0
Soybean, hulls	5.0

(b) Section 18 emergency exemptions. [Reserved]

Environmental Protection Agency

§ 180.474

(c) *Tolerances with regional restrictions.* [Reserved]

(d) *Indirect or inadvertent residues.* Tolerances are established for indirect or inadvertent residues of the herbicide glufosinate ammonium, butanoic acid, 2-amino-4-(hydroxymethylphosphinyl)-, monoammonium salt and its metabolite, 3-methylphosphinopropionic acid in or on the following raw agricultural commodities when present therein as a result of the application of glufosinate ammonium to crops listed in paragraph (a) of this section:

Commodity	Parts per million
Barley, hay	0.40
Barley, straw	0.40
Buckwheat, fodder	0.40
Buckwheat, forage	0.40
Oat, forage	0.40
Oat, hay	0.40
Oat, straw	0.40
Rye, forage	0.40
Rye, straw	0.40
Teosinte	0.40
Triticale	0.40
Wheat, forage	0.40
Wheat, hay	0.40
Wheat, straw	0.40

[68 FR 55849, Sept. 29, 2003, as amended at 71 FR 25945, May 3, 2006; 72 FR 72625, Dec. 21, 2007]

§ 180.474 Tebuconazole; tolerances for residues.

(a) *General.* (1) Tolerances are established for residues of the fungicide tebuconazole, including its metabolites and degradates, in or on the commodities in the following table. Compliance with the tolerance levels specified in the following table is to be determined by measuring only tebuconazole (alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol), in or on the commodity.

Commodity	Parts per million
Almond, hulls	6.0
Apple, wet pomace	0.1
Asparagus	0.05
Banana	0.05
Barley, grain	0.15
Barley, hay	7.0
Barley, straw	3.5
Bean, dry seed	0.1
Bean, succulent	0.1
Beet, garden, roots	0.70
Beet, garden, tops	7.0
Brassica, leafy greens, subgroup 5B	2.5
Cherry, sweet, pre- and post-harvest	5.0
Cherry, tart, pre- and post-harvest	5.0

Commodity	Parts per million
Coffee, green bean ¹	0.15
Coffee, roasted bean ¹	0.3
Corn, field, forage	4.0
Corn, field, grain	0.05
Corn, field, stover	3.5
Corn, pop, grain	0.05
Corn, pop, stover	3.5
Corn, sweet, forage	7.0
Corn, sweet, kernel plus cob with husks removed	0.5
Corn, sweet, stover	6.0
Cotton, gin byproducts	25.0
Cotton, undelinted seed	2.0
Fruit, pome, group 11	0.05
Fruit, stone, group 12, except cherry	1.0
Grain, aspirated fractions	16.0
Grape	5.0
Grass, forage	8.0
Grass, hay	25.0
Grass, seed screenings	55.0
Grass, straw	30.0
Hop, dried cones	35.0
Lychee	1.6
Mango, postharvest	0.15
Nut, tree, group 14	0.05
Oat, forage	0.10
Oat, grain	0.05
Oat, hay	0.10
Oat, straw	0.10
Okra	1.2
Onion, bulb, subgroup 3-07A	0.2
Onion, green, subgroup 3-07B	1.3
Peach	1.0
Peanut	0.1
Pistachio	0.05
Plum, pre- and post-harvest	1.0
Soybean, forage	25
Soybean, hay	50
Soybean, seed	0.08
Sunflower, seed	0.05
Sunflower, meal	0.2
Sunflower, refined oil	0.2
Vegetable, cucurbit, group 9	0.09
Vegetable, fruiting, group 8	1.3
Wheat, forage	3.0
Wheat, grain	0.05
Wheat, hay	7.0
Wheat, straw	1.5

¹There are no U.S. registrations as of 7/31/2008.

(2) Tolerances are established for residues of the fungicide tebuconazole, including its metabolites and degradates, in or on the commodities in the following table. Compliance with the tolerance levels specified in the following table is to be determined by measuring only the sum of tebuconazole (alpha-[2-(4-chlorophenyl)ethyl]-alpha-(1,1-dimethylethyl)-1H-1,2,4-triazole-1-ethanol) and its diol metabolite (1-(4-chlorophenyl)-4,4-dimethyl-3-(1H-1,2,4-triazole-1-yl-methyl)-pentane-3,5-diol), calculated as the stoichiometric equivalent of tebuconazole, in or on the commodity.