

§ 320.23

§ 320.23 Basis for measuring in vivo bioavailability or demonstrating bioequivalence.

(a)(1) The in vivo bioavailability of a drug product is measured if the product's rate and extent of absorption, as determined by comparison of measured parameters, e.g., concentration of the active drug ingredient in the blood, urinary excretion rates, or pharmacological effects, do not indicate a significant difference from the reference material's rate and extent of absorption. For drug products that are not intended to be absorbed into the bloodstream, bioavailability may be assessed by measurements intended to reflect the rate and extent to which the active ingredient or active moiety becomes available at the site of action.

(2) Statistical techniques used shall be of sufficient sensitivity to detect differences in rate and extent of absorption that are not attributable to subject variability.

(3) A drug product that differs from the reference material in its rate of absorption, but not in its extent of absorption, may be considered to be bioavailable if the difference in the rate of absorption is intentional, is appropriately reflected in the labeling, is not essential to the attainment of effective body drug concentrations on chronic use, and is considered medically insignificant for the drug product.

(b) Two drug products will be considered bioequivalent drug products if they are pharmaceutical equivalents or pharmaceutical alternatives whose rate and extent of absorption do not show a significant difference when administered at the same molar dose of the active moiety under similar experimental conditions, either single dose or multiple dose. Some pharmaceutical equivalents or pharmaceutical alternatives may be equivalent in the extent of their absorption but not in their rate of absorption and yet may be considered bioequivalent because such differences in the rate of absorption are intentional and are reflected in the labeling, are not essential to the attainment of effective body drug concentrations on chronic use, and are consid-

21 CFR Ch. I (4-1-11 Edition)

ered medically insignificant for the particular drug product studied.

[57 FR 17999, Apr. 28, 1992, as amended at 67 FR 77673, Dec. 19, 2002]

§ 320.24 Types of evidence to measure bioavailability or establish bioequivalence.

(a) Bioavailability may be measured or bioequivalence may be demonstrated by several in vivo and in vitro methods. FDA may require in vivo or in vitro testing, or both, to measure the bioavailability of a drug product or establish the bioequivalence of specific drug products. Information on bioequivalence requirements for specific products is included in the current edition of FDA's publication "Approved Drug Products with Therapeutic Equivalence Evaluations" and any current supplement to the publication. The selection of the method used to meet an in vivo or in vitro testing requirement depends upon the purpose of the study, the analytical methods available, and the nature of the drug product. Applicants shall conduct bioavailability and bioequivalence testing using the most accurate, sensitive, and reproducible approach available among those set forth in paragraph (b) of this section. The method used must be capable of measuring bioavailability or establishing bioequivalence, as appropriate, for the product being tested.

(b) The following in vivo and in vitro approaches, in descending order of accuracy, sensitivity, and reproducibility, are acceptable for determining the bioavailability or bioequivalence of a drug product.

(1)(i) An in vivo test in humans in which the concentration of the active ingredient or active moiety, and, when appropriate, its active metabolite(s), in whole blood, plasma, serum, or other appropriate biological fluid is measured as a function of time. This approach is particularly applicable to dosage forms intended to deliver the active moiety to the bloodstream for systemic distribution within the body; or

(ii) An in vitro test that has been correlated with and is predictive of human in vivo bioavailability data; or