

labratus, *Cyathostomum* spp. including *C. catinatum* and *C. pateratum*, *Cylicocyclus* spp. including *C. insigne*, *C. leptostomum*, *C. nassatus*, and *C. brevicapsulatus*, *Cylicodontophorus* spp., *Cylicostephanus* spp. including *C. calicatus*, *C. goldi*, *C. longibursatus*, and *C. minutus*, and *Petrovinema poculatum*; Small Strongyles—fourth-stage larvae; Pinworms (adults and fourth-stage larvae)—*Oxyuris equi*; Ascarids (adults and third- and fourth-stage larvae)—*Parascaris equorum*; Hairworms (adults)—*Trichostrongylus axei*; Large-mouth Stomach Worms (adults)—*Habronema muscae*; Bots (oral and gastric stages)—*Gasterophilus* spp. including *G. intestinalis* and *G. nasalis*; Lungworms (adults and fourth-stage larvae)—*Dictyocaulus arnfieldi*; Intestinal Threadworms (adults)—*Strongyloides westeri*; Summer Sores caused by *Habronema* and *Draschia* spp. cutaneous third-stage larvae; Dermatitis caused by neck threadworm microfilariae, *Onchocerca* sp.

(ii) Tapeworms—*Anoplocephala perfoliata*; Large Strongyles (adults)—*Strongylus vulgaris* (also early forms in blood vessels), *S.edentatus* (also tissue stages), *S. equinus*, *Triodontophorus* spp.; Small Strongyles (adults, including those resistant to some benzimidazole class compounds)—*Cyathostomum* spp., *Cylicocyclus* spp., *Cylicostephanus* spp., *Cylicodontophorus* spp.; Small Strongyles—fourth-stage larvae; Pinworms (adults and fourth-stage larvae)—*Oxyuris equi*; Ascarids (adults and third- and fourth-stage larvae)—*Parascaris equorum*; Hairworms (adults)—*Trichostrongylus axei*; Large-mouth Stomach Worms (adults)—*Habronema muscae*; Bots (oral and gastric stages)—*Gasterophilus* spp.; Lungworms (adults and fourth-stage larvae)—*Dictyocaulus arnfieldi*; Intestinal Threadworms (adults)—*Strongyloides westeri*; Summer Sores caused by *Habronema* and *Draschia* spp. cutaneous third-stage larvae; Dermatitis caused by neck threadworm microfilariae, *Onchocerca* sp.

(3) *Limitations*. For oral use only. Do not use in horses intended for human consumption.

[68 FR 55309, Sept. 25, 2003, as amended at 69 FR 49808, Aug. 12, 2004; 70 FR 65835, Nov. 1, 2005]

§520.1199 Ivermectin, pyrantel, and praziquantel tablets.

(a) *Specifications*. Each chewable tablet contains:

(1) 34 micrograms (mcg) ivermectin, 28.5 milligrams (mg) pyrantel pamoate, and 28.5 mg praziquantel;

(2) 68 mcg ivermectin, 57 mg pyrantel pamoate, and 57 mg praziquantel;

(3) 136 mcg ivermectin, 114 mg pyrantel pamoate, and 114 mg praziquantel; or

(4) 272 mcg ivermectin, 228 mg pyrantel pamoate, and 228 mg praziquantel.

(b) *Sponsors*. See No. 051311 in §510.600(c) of this chapter.

(c) *Conditions of use in dogs*—(1) *Amount*. Administer monthly according to body weight as follows:

(i) 6 to 12 lb: one tablet as described in paragraph (a)(1) of this section.

(ii) 12.1 to 25 lb: one tablet as described in paragraph (a)(2) of this section.

(iii) 25.1 to 50 lb: one tablet as described in paragraph (a)(3) of this section.

(iv) 50.1 to 100 lb: one tablet as described in paragraph (a)(4) of this section.

(v) Greater than 100 lb: use the appropriate combination of tablets.

(2) *Indications for use*. Prevents canine heartworm disease by eliminating the tissue stage of heartworm larvae (*Dirofilaria immitis*) for 1 month (30 days) after infection and for the treatment and control of roundworm (*Toxocara canis*, *Toxascaris leonina*), hookworm (*Ancylostoma caninum*, *Uncinaria stenocephala*, *Ancylostoma braziliense*) and tapeworm (*Dipylidium caninum*, *Taenia pisiformis*) infections.

(3) *Limitations*. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[71 FR 65052, Nov. 7, 2006]

§520.1200 Ivermectin, fenbendazole, and praziquantel tablets.

(a) *Specifications*. Each chewable tablet contains either:

(1) 68 micrograms (µg) ivermectin, 1.134 grams fenbendazole, and 57 milligrams (mg) praziquantel; or

(2) 27 µg ivermectin, 454 mg fenbendazole, and 23 mg praziquantel.