### Food and Drug Administration, HHS

520.2520f Trichlorofon granules.

520.2520g Trichlorfon, phenothiazine, and piperazine dihydrochloride powder.

520.2582 Triflupromazine hydrochloride tablets.

520.2598 Trilostane.

520.2604 Trimeprazine tartrate and prednisolone tablets.

520.2605 Trimeprazine tartrate and prednisolone capsules.

520.2610 Trimethoprim and sulfadiazine tablets.

520.2611 Trimethoprim and sulfadiazine paste.

520.2612 Trimethoprim and sulfadiazine oral suspension.

520.2613 Trimethoprim and sulfadiazine powder.

520.2640 Tylosin.

AUTHORITY: 21 U.S.C. 360b.

SOURCE: 40 FR 13838, Mar. 27, 1975, unless otherwise noted.

#### § 520.23 Acepromazine.

(a) *Specifications*. Each tablet contains 5, 10, or 25 milligrams (mg) acepromazine maleate.

(b) Sponsors. See No. 000010 in §510.600(c) of this chapter.

(c) Conditions of use—(1) Dogs—(i) Amount. 0.25 to 1.0 mg per pound (/lb) body weight orally.

(ii) Indications for use. As an aid in tranquilization and as a preanesthetic agent.

(iii) *Limitations*. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) Cats—(i) Amount. 0.5 to 1.0 mg/lb body weight orally.

(ii) *Indications for use*. As a tranquilizer.

(iii) *Limitations*. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[75 FR 10165, Mar. 5, 2010]

## §520.44 Acetazolamide sodium soluble powder.

(a) *Specifications*. The drug is in a powder form containing acetazolamide sodium, USP equivalent to 25 percent acetazolamide activity.

(b) Sponsor. See No. 053501 in \$510.600(c) of this chapter.

(c) Conditions of use. (1) It is used in dogs as an aid in the treatment of mild

congestive heart failure and for rapid reduction of intraocular pressure.<sup>1</sup>

(2) It is administered orally at a dosage level of 5 to 15 milligrams per pound of body weight daily.<sup>1</sup>

(3) For use only by or on the order of a licensed veterinarian.<sup>1</sup>

 $[40\ {\rm FR}\ 13838,\ {\rm Mar.}\ 27,\ 1975,\ {\rm as}\ {\rm amended}\ {\rm at}\ 67\ {\rm FR}\ 78355,\ {\rm Dec.}\ 24,\ 2002]$ 

# § 520.45 Albendazole oral dosage forms.

#### §520.45a Albendazole suspension.

(a) *Specifications*. Each milliliter of suspension contains 45.5 milligrams (mg) (4.55 percent) or 113.6 mg (11.36 percent) albendazole.

(b) Sponsor. See No. 000069 in §510.600 of this chapter.

(c) *Related tolerances*. See §556.34 of this chapter.

(d) *Special considerations*. See §500.25 of this chapter.

(e) Conditions of use—(1) Cattle. Administer 11.36 percent suspension:

(i) Amount. 4.54 mg/pound (lb) body weight (10 mg/kilogram (kg)) as a single oral dose using dosing gun or dosing syringe.

(ii) Indications for use. For removal and control of adult liver flukes (Fasciola hepatica); heads and segments of tapeworms (Moniezia benedeni and M. expanse); adult and 4th stage larvae of stomach worms (brown stomach worms including 4th stage inhibited larvae (Ostertagia ostertagi), barberpole worm (Haemonchus contortus and H. placei), small stomach worm (Trichostrongylus axei)); adult and 4th stage larvae of intestinal worms (thread-necked intestinal worm (Nematodirus spathiger and N. helvetianus), small intestinal worm (Cooperia punctata and C. oncophora)); adult stages of intestinal worms (hookworm (Bunostomum phlebotomum), bankrupt worm (Trichostrongylus colubriformis), nodular worm (Oesophagostomum radiatum)); adult and 4th stage larvae of lungworms (Dictyocaulus viviparus).

<sup>1</sup>These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by §514.111 of this chapter, but may require bioequivalency and safety information.

## §520.45a