assistance in the diagnosis, treatment, and control of parasitism.

[50 FR 10221, Mar. 14, 1985, as amended at 62 FR 63270, Nov. 28, 1997]

§520.522 Cyclosporine.

(a) *Specifications*. Each capsule contains 10, 25, 50, or 100 milligrams (mg) cyclosporine.

(b) Sponsor. See No. 058198 in \$510.600(c) of this chapter.

(c) [Reserved]

(d) Conditions of use in dogs—(1) Amount. 5 mg per kilogram of body weight given orally as a single daily dose for 30 days. Following this initial daily treatment period, the dosage may be tapered by decreasing the frequency of administration to every other day or two times a week, until a minimum frequency is reached which will maintain the desired therapeutic effect.

(2) *Indications for use*. For the control of atopic dermatitis in dogs weighing at least 4 pounds body weight.

(3) *Limitations*. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[68 FR 54804, Sept. 19, 2003]

§ 520.530 Cythioate oral liquid.

(a) *Specifications*. Each milliliter contains 15 milligrams of cythioate.

(b) *Sponsor*. See Nos. 000859 and 053501 in §510.600 of this chapter.

(c) Special considerations. Cythioate is a cholinesterase inhibitor. Do not use this product in animals simultaneously with or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, insecticides, pesticides, or chemicals.

(d) Conditions of use—(1) Amount. 15 milligrams cythicate per 10 pounds of body weight every third day or twice a week.

(2) *Indications for use*. Dogs, for control of fleas.

(3) *Limitations*. For oral use in dogs only. Do not use in greyhounds or in animals that are pregnant, sick, under stress, or recovering from surgery. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

 $[49\ {\rm FR}\ 5614,\ {\rm Feb}.\ 14,\ 1984,\ {\rm as}\ {\rm amended}\ {\rm at}\ 67\ {\rm FR}\ 78355,\ {\rm Dec.}\ 24,\ 2002]$

21 CFR Ch. I (4–1–11 Edition)

§ 520.531 Cythioate tablets.

(a) [Reserved]

(b) *Sponsors*. See No. 000859 in §510.600(c) of this chapter for use of 30and 90-milligram (mg) tablets and see No. 053501 in §510.600(c) of this chapter for use of 30-mg tablet.

(c) Special considerations. Cythioate is a cholinesterase inhibitor. Do not use this product in animals simultaneously with or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, insecticides, pesticides, or chemicals.

(d) Conditions of use—(1) Amount. 30 milligrams cythicate per 20 pounds of body weight every third day or twice a week.

(2) *Indications for use*. Dogs, for control of fleas.

(3) *Limitations.* For oral use in dogs only. Do not use in greyhounds or in animals that are pregnant, sick, under stress, or recovering from surgery. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[49 FR 5615, Feb. 14, 1984, as amended at 59 FR 26942, May 25, 1994; 67 FR 78355, Dec. 24, 2002]

§ 520.534 Decoquinate.

(a) Specifications. The drug is a powder containing 0.8 percent decoquinate.

(b) Sponsor. See No. 046573 in §510.600(c) of this chapter.

(c) *Related tolerances*. See §556.170 of this chapter.

(d) Conditions of use. Calves—(1) Amount. Feed 22.7 milligrams per 100 pounds of body weight (0.5 milligram per kilogram) per day.

(2) Indications for use. For the prevention of coccidiosis in ruminating and nonruminating calves, including veal calves, caused by *Eimeria bovis* and *E.* zuernii.

(3) *Limitations.* Feed in whole milk at the rate of 22.7 milligrams per 100 pounds body weight daily (0.5 milligram per kilogram) for at least 28 days.

[64 FR 10103, Mar. 2, 1999, as amended at 64 FR 30386, June 8, 1999]

§520.538 Deracoxib.

(a) *Specifications*. Each chewable tablet contains 25, 50, 75, or 100 milligrams (mg) deracoxib.