

§ 522.723

21 CFR Ch. I (4-1-11 Edition)

ovulation in estrous cycling cattle that have a corpus luteum.

(ii) *Beef cattle and nonlactating dairy heifers*—(A) *Amount.* 25 mg as a single intramuscular injection.

(B) *Indications.* For treatment of pyometra (chronic endometritis).

(iii) *Nonlactating cattle*—(A) *Amount.* 25 mg as a single intramuscular injection during the first 100 days of gestation.

(B) *Indications.* For its abortifacient effect in nonlactating cattle.

(iv) *Lactating dairy cattle*—(A) *Amount.* 25 mg as a single intramuscular injection.

(B) *Indications.* For treatment of unobserved (silent) estrus in lactating dairy cattle that have a corpus luteum.

(v) Dinoprost solution as provided by No. 000009 in §510.600(c) of this chapter may be used concurrently with progesterone intravaginal inserts as in §529.1940 of this chapter.

(3) *Swine*—(i) *Amount.* 10 mg as a single intramuscular injection.

(ii) *Indications.* For parturition induction in swine when injected within 3 days of normal predicted farrowing.

[67 FR 41824, June 20, 2002]

§ 522.723 Diprenorphine hydrochloride injection.

(a) *Chemical name.* N-(Cyclopropylmethyl)-6,7,8,14-tetrahydro-7- α -(1-hydroxy - 1 - methylethyl) - 6,14 - endoethanonoripavine hydrochloride.

(b) *Specifications.* Each milliliter of diprenorphine hydrochloride injection, veterinary, contains 2 mg of diprenorphine hydrochloride in sterile aqueous solution.

(c) *Sponsors.* See No. 053923 in §510.600(c) of this chapter.

(d) *Conditions of use.* (1) The drug is used for reversing the effects of etorphine hydrochloride injection, veterinary, the use of which is provided for in §522.883, in wild and exotic animals.

(2) It is administered intramuscularly or intravenously at a suitable dosage level depending upon the species.

(3) For use in wild or exotic animals only. Do not use in domestic food-producing animals. Do not use 30 days before, or during, the hunting season in

free-ranging wild animals that might be used for food.

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian. Distribution is restricted to veterinarians engaged in zoo and exotic animal practice, wildlife management programs and researchers.

[40 FR 13858, Mar. 27, 1975, as amended at 48 FR 16241, Apr. 15, 1983; 60 FR 39847, Aug. 4, 1995; 64 FR 15684, Apr. 1, 1999]

§ 522.770 Doramectin.

(a) *Specifications.* Each milliliter of sterile aqueous solution contains 10 milligrams of doramectin.

(b) *Sponsor.* See No. 000069 in §510.600(c) of this chapter.

(c) *Related tolerances.* See §556.225 of this chapter.

(d) *Conditions of use*—(1) *Cattle*—(i) *Amount.* 200 micrograms per kilogram (10 milligrams per 110 pounds).

(ii) *Indications for use.* For treatment and control of gastrointestinal roundworms, lungworms, eyeworms, grubs, sucking lice, and mange mites. To control infections and to protect from reinfection with *Cooperia oncophora* and *Haemonchus placei* for 14 days, *Ostertagia ostertagi* for 21 days, and *C. punctata*, *Oesophagostomum radiatum*, and *Dictyocaulus viviparus* for 28 days after treatment.

(iii) *Limitations.* Administer as a single subcutaneous or intramuscular injection. Do not slaughter cattle within 35 days of treatment. Not for use in female dairy cattle 20 months of age or older. Do not use in calves to be processed for veal.

(2) *Swine*—(i) *Amount.* 300 micrograms per kilogram (10 milligrams per 75 pounds).

(ii) *Indications for use.* For treatment and control of gastrointestinal roundworms, lungworms, kidney worms, sucking lice, and mange mites.

(iii) *Limitations.* Administer as a single intramuscular injection. Do not slaughter swine within 24 days of treatment. Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[61 FR 53321, Oct. 11, 1996, as amended at 62 FR 44410, Aug. 21, 1997; 62 FR 62242, Nov. 21, 1997; 63 FR 68183, Dec. 10, 1998; 64 FR 13509, Mar. 19, 1999]