

once or twice daily, and repeat treatment as required. Use only as recommended by a veterinarian in treatment of puncture wounds, wounds requiring surgical debridement or suturing, those of a chronic nature involving proud flesh, generalized and chronic infections of the skin, and those skin conditions associated with intense itching. If redness, irritation, or swelling persists or increases, discontinue use and consult a veterinarian. Not for use in horses intended for food.

[45 FR 49543, July 25, 1980, as amended at 50 FR 30153, July 24, 1985; 56 FR 50653, Oct. 8, 1991; 57 FR 31314, July 15, 1992; 60 FR 55659, Nov. 2, 1995; 65 FR 41588, July 6, 2000]

§ 524.1044 Gentamicin sulfate ophthalmic and topical dosage forms.

§ 524.1044a Gentamicin ophthalmic solution.

(a) *Specifications.* Each milliliter of sterile aqueous solution contains gentamicin sulfate equivalent to 3 milligrams of gentamicin.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used in dogs and cats for the topical treatment of infections of the conjunctiva caused by susceptible bacteria.

(2) Administer 1 or 2 drops into the conjunctival sac 2 to 4 times a day.

(3) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[41 FR 14189, Apr. 2, 1976, as amended at 52 FR 7832, Mar. 13, 1987]

§ 524.1044b Gentamicin sulfate, betamethasone valerate otic solution.

(a) *Specifications.* Each milliliter of solution contains gentamicin sulfate equivalent to 3 milligrams (mg) gentamicin base and betamethasone valerate equivalent to 1 mg betamethasone alcohol.

(b) *Sponsors.* See Nos. 000061 and 054925 in § 510.600(c) of this chapter.

(c) *Conditions of use—*(1) *Amounts and indications for use—*(i) For the treatment of acute and chronic otitis externa caused by bacteria sensitive to gentamicin in dogs, instill three to

eight drops of solution into the ear canal twice daily for 7 to 14 days.

(ii) For the treatment of infected superficial lesions caused by bacteria sensitive to gentamicin in dogs and cats, apply a sufficient amount of the drug to cover the treatment area twice daily for 7 to 14 days.

(2) *Limitations.* Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[71 FR 13542, Mar. 16, 2006]

§ 524.1044c Gentamicin sulfate ophthalmic ointment.

(a) *Specifications.* Each gram of ointment contains gentamicin sulfate equivalent to 3 milligrams of gentamicin.

(b) *Sponsors.* See Nos. 000061 and 025463 in § 510.600(c) of this chapter.

(c) *Conditions of use in dogs and cats—*

(1) *Amount.* Apply approximately a 1/2-inch strip to the affected eye 2 to 4 times a day.

(2) *Indications for use.* For treatment of conjunctivitis caused by susceptible bacteria.

(3) *Limitations.* Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[69 FR 47363, Aug. 5, 2004]

§ 524.1044d Gentamicin sulfate, betamethasone valerate ointment.

(a) *Specifications.* Each gram of ointment contains gentamicin sulfate equivalent to 3 milligrams of gentamicin base and betamethasone valerate equivalent to 1 milligram of betamethasone.

(b) *Sponsor.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Conditions of use.* (1) The drug is used or indicated for use in dogs in the treatment of acute and chronic canine otitis externa and canine infected superficial lesions caused by bacteria sensitive to gentamicin.

(2)(i) For the treatment of acute and chronic canine otitis externa the drug is administered by instillation of 3 to 8 drops into the ear canal twice daily for 7 days. The external ear and ear canal should be properly cleaned and dried before treatment. Remove foreign material, debris, crusted exudates, etc., with suitable nonirritating solutions.