

ceftiofur is 0.830 milligrams per kilogram of body weight. The ASDI is the amount of total residues of ceftiofur that may safely be consumed in a single meal. The ASDI is used to derive the tolerance for residues of desfuoylceftiofur at the injection site.

(b) *Tolerances*—(1) *Poultry, and sheep*. A tolerance for residues of ceftiofur in edible tissue is not required.

(2) *Swine*. The tolerances for desfuoylceftiofur (marker residue) are:

(i) *Kidney (target tissue)*. 0.25 parts per million (ppm).

(ii) *Liver*. 3 ppm.

(iii) *Muscle*. 2 ppm.

(3) *Cattle*. The tolerances for desfuoylceftiofur (marker residue) are:

(i) *Kidney (target tissue)*. 0.4 ppm.

(ii) *Liver*. 2 ppm.

(iii) *Muscle*. 1 ppm.

(iv) *Milk*. 0.1 ppm.

[63 FR 53579, Oct. 6, 1998, as amended at 68 FR 60296, Oct. 22, 2003; 69 FR 43892, July 23, 2004; 71 FR 39546, July 13, 2006]

#### § 556.115 Cephapirin.

A tolerance of 0.02 parts per million (ppm) is established for residues of cephalapirin in the milk and 0.1 ppm in the uncooked edible tissues of dairy cattle.

[40 FR 57454, Dec. 10, 1975]

#### § 556.120 Chlorhexidine.

A tolerance of zero is established for residues of chlorhexidine in the uncooked edible tissues of calves.

#### § 556.150 Chlortetracycline.

(a) *Acceptable daily intake (ADI)*. The ADI for total residues of tetracyclines including chlortetracycline, oxytetracycline, and tetracycline is 25 micrograms per kilogram of body weight per day.

(b) *Tolerances*. (1) Tolerances are established for the sum of tetracycline residues in tissues of beef cattle, non-lactating dairy cows, calves, swine, sheep, chickens, turkeys, and ducks, of 2 parts per million (ppm) in muscle, 6 ppm in liver, and 12 ppm in fat and kidney.

(2) A tolerance is established for residues of chlortetracycline in eggs of 0.4 ppm.

[63 FR 52158, Sept. 30, 1998, as amended at 63 FR 57246, Oct. 27, 1998]

#### § 556.160 Clopidol.

Tolerances for residues of clopidol (3,5-dichloro-2,6-dimethyl-4-pyridinol) in food are established as follows:

(a) In cereal grains, vegetables, and fruits: 0.2 part per million.

(b) In chickens and turkeys:

(1) 15 parts per million in uncooked liver and kidney.

(2) 5 parts per million in uncooked muscle.

(c) In cattle, sheep, and goats:

(1) 3 parts per million in uncooked kidney.

(2) 1.5 parts per million in uncooked liver.

(3) 0.2 part per million in uncooked muscle.

(d) In swine: 0.2 part per million in uncooked edible tissues.

(e) In milk: 0.02 part per million (negligible residue).

#### § 556.163 Clorsulon.

(a) *Acceptable daily intake (ADI)*. The ADI for total residues of clorsulon is 8 micrograms per kilogram of body weight per day.

(b) *Tolerances*—(1) *Cattle*—(i) *Kidney (the target tissue)*. The tolerance for parent clorsulon (the marker residue) is 1.0 part per million.

(ii) *Muscle*. The tolerance for parent clorsulon (the marker residue) is 0.1 part per million.

(2) [Reserved]

[66 FR 35544, July 6, 2001]

#### § 556.165 Cloxacillin.

A tolerance of 0.01 part per million is established for negligible residues of cloxacillin in the uncooked edible tissues of cattle and in milk.

[40 FR 28792, July 9, 1975]

#### § 556.167 Colistimethate.

A tolerance for residues of colistimethate in the edible tissues of chickens is not required.

[63 FR 13123, Mar. 18, 1998]