

§ 558.248

21 CFR Ch. I (4–1–11 Edition)

(B) *Limitations.* Feed continuously as sole ration. Not to be used in swine weighing more than 250 pounds.

(ii) *Amount.* 3.6 to 14.5 grams per ton.

(A) *Indications for use.* For increased rate of weight gain.

(B) *Limitations.* Feed continuously as sole ration. Not to be used in swine weighing more than 250 pounds.

(2) [Reserved]

[57 FR 38442, Aug. 25, 1992, as amended at 62 FR 63271, Nov. 28, 1997]

§ 558.248 Erythromycin thiocyanate.

(a) *Approvals.* Type A medicated articles: (1) 2.2 percent to 061623 in

§ 510.600(c) of this chapter for use as in paragraph (d) of this section.

(2) 5 and 10 percent to 061623 for use in paragraphs (d)(1)(i) and (ii) of this section.

(b) *Special considerations.* The levels of antibiotic are expressed in terms of erythromycin master standard. One gram of erythromycin thiocyanate is equivalent to 0.925 gram of erythromycin master standard.

(c) *Related tolerances.* See § 556.230 of this chapter.

(d) *Condition of use.* (1) It is used as follows:

| Erythromycin thiocyanate in grams per ton | Combination in grams per ton | Indications for use | Limitations | Sponsor |
|-------------------------------------------|------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------|
| (i) 4.6 to 18.5 | | Chickens; growth promotion and feed efficiency. | | 061623 |
| (ii) 9.25 to 18.5 ... | | Turkeys; growth promotion and feed efficiency. | For turkeys not over 12 weeks of age. | 061623 |
| (iii) 9.25 to 64.75 | | Swine; increase in weight gain, improved feed efficiency in starter pigs (9.25 to 64.75) and grower-finishing pigs (9.25). | Starter ration for animals up to 35 lb body weight. | 061623 |
| (iv) 18.5 | | Laying chickens; aids in increasing egg production. | | 061623 |
| (v) 92.5 | | 1. Chickens; as an aid in the prevention of chronic respiratory disease during periods of stress. 2. Chickens; as an aid in the prevention of infectious coryza. 3. Turkeys; as an aid in the prevention of chronic respiratory disease during periods of stress. | Feed for 2 d before stress and 3 to 6 d after stress; withdraw 24 h before slaughter. Feed for 7 to 14 d; withdraw 24 h before slaughter. Feed for 2 d before stress and 3 to 6 d after stress. | 061623 |
| (vi) 185 | | 1. Chickens; as an aid in the prevention and reduction of lesions and in lowering severity of chronic respiratory disease. 2. Turkeys; as an aid in the prevention and reduction of lesions and in lowering severity of chronic respiratory disease. | Feed for 5 to 8 d; do not use in birds producing eggs for food purposes; withdraw 48 h before slaughter. Feed for 5 to 8 d; do not use in birds producing eggs for food purposes. | 061623 061623 |

(2) In feed for feedlot beef cattle at 37 milligrams per head per day as an aid in stimulating growth and improving feed efficiency.

(3) Erythromycin thiocyanate may be used in accordance with the provisions of this section in the combinations provided as follows:

(i) Amprolium in accordance with § 558.55.

(ii) Amprolium and ethopabate in accordance with § 558.58.

(iii) Arsanilic acid in accordance with § 558.62.

(iv) Zoalene in accordance with § 558.680.

[41 FR 10999, Mar. 15, 1976, as amended at 45 FR 56799, Aug. 26, 1980; 49 FR 31281, Aug. 6, 1984; 51 FR 7397, Mar. 3, 1986; 52 FR 2684, Jan. 26, 1987; 54 FR 12189, Mar. 24, 1989; 66 FR 14074, Mar. 9, 2001; 68 FR 4915, Jan. 31, 2003]

§ 558.254 Famphur.

(a) *Approvals.* Type A medicated articles: 13.2 and 33.3 percent to 000061 in § 510.600(c) of this chapter.

(b) *Special considerations.* Famphur is a cholinesterase inhibitor. Do not use this product in animals simultaneously or within a few days before or after

Food and Drug Administration, HHS

§ 558.258

treatment with or exposure to cholinesterase-inhibiting drugs, pesticides, or chemicals.

(c) *Related tolerances.* See § 556.273 of this chapter.

(d) *Conditions of use.* It is used in the feed for cattle as follows:

(1) *Amount.* 1.1 milligrams per pound body weight per day.

(i) *Indications for use.* For control of grubs and as an aid in control of sucking lice.

(ii) *Limitations.* For beef cattle and nonlactating dairy cows; feed for 30 days; withdraw from dry dairy cows and heifers 21 days prior to freshening; withdraw 4 days prior to slaughter.

(2) *Amount.* 2.3 milligrams per pound body weight per day.

(i) *Indications for use.* For control of grubs.

(ii) *Limitations.* For beef cattle and nonlactating dairy cows; feed for 10 days; withdraw from dry dairy cows and heifers 21 days prior to freshening; withdraw 4 days prior to slaughter.

[41 FR 11000, Mar. 15, 1976, as amended at 51 FR 7397, Mar. 3, 1986; 57 FR 7652, Mar. 4, 1992; 62 FR 55161, Oct. 23, 1997; 62 FR 61626, Nov. 19, 1997]

§ 558.258 Fenbendazole.

(a) *Specifications.* Type A medicated articles: 4 percent (18.1 grams per pound (g/lb)), 8 percent (36.2 g/lb), and 20 percent (90.7 g/lb) fenbendazole.

(b) *Approvals.* See No. 000061 in § 510.600(c) of this chapter.

(c) *Related tolerances.* See § 556.275 of this chapter.

(d) *Special considerations.* See § 500.25 of this chapter.

(e) *Conditions of use—(1) Turkeys.*

| Amount fenbendazole in grams per ton | Combination in grams per ton | Indications for use | Limitations | Sponsor |
|--------------------------------------|------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------|---------|
| 14.5 (16 parts per million). | | Growing turkeys: For the removal and control of gastrointestinal worms: roundworms, adult and larvae (<i>Ascaridia dissimilis</i>); cecal worms, adult and larvae (<i>Heterakis gallinarum</i>), an important vector of <i>Histomonas meleagridis</i> (Black-head). | Feed continuously as the sole ration for 6 days. For growing turkeys only. | 000061 |

(2) *Swine.*

| Amount fenbendazole in grams per ton | Combination in grams per ton | Indications for use | Limitations | Sponsor |
|---------------------------------------------------------------------------------------------------------------|------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------|---------|
| (i) 10 to 300 (to provide 9 milligrams per kilogram (mg/kg) of body weight) given over a 3- to 12-day period. | | For the removal and control of: Adult stage lungworms (<i>Metastrongylus apri</i> and <i>M. pudendotectus</i>); adult and larvae (L3, 4 stages—liver, lung, intestinal forms) large roundworms (<i>Ascaris suum</i>); adult stage nodular worms (<i>Oesophagostomum dentatum</i> , <i>O. quadrispinulatum</i>); adult stage small stomach worms (<i>Hyostromylus rubidus</i>); adult and larvae (L2, 3, 4 stages—intestinal mucosal forms) whipworms (<i>Trichuris suis</i>); adult and larvae kidney worms (<i>Stephanurus dentatus</i>). | Feed as sole ration | 000061 |
| (ii) 10 to 80 (to provide 9 mg/kg of body weight). | Lincomycin 20 | As in paragraph (e)(2)(i) of this section; for increased rate of gain in growing-finishing swine. | Feed as sole ration. Do not feed to swine that weigh more than 250 pounds (lbs); lincomycin as provided by 000009 in § 510.600(c) of this chapter. | 000061 |