

§ 573.940

EFFECTIVE DATE NOTE: At 58 FR 47973, Sept. 13, 1993, the amendments to § 573.920 that were published at 52 FR 10887, Apr. 6, 1987; 52 FR 21001, June 4, 1987; and 54 FR 14214, Apr. 10, 1989 were stayed until further notice. At 59 FR 45973, Sept. 6, 1994, the stay was confirmed.

§ 573.940 Silicon dioxide.

The food additive silicon dioxide may be safely used in animal feed in accordance with the following conditions:

(a) The food additive is manufactured by vapor phase hydrolysis or by other means whereby the particle size is such as to accomplish the intended effect.

(b) It is used or intended for use in feed components as an anticaking agent, and/or grinding aid, as follows:

Feed component	Limitations (percent)
BHT (butylated hydroxytoluene)	2
Methionine hydroxy analog and its calcium salts	1
Piperazine, piperazine salts	0.8
Sodium propionate	1
Urea	1
Vitamins	3

(c) It is used in feed as an anticaking agent in an amount not to exceed that reasonably required to accomplish its intended effect and in no case in an amount to exceed 2 percent by weight of the finished feed.

§ 573.960 Sorbitan monostearate.

The food additive sorbitan monostearate may be safely used alone or in combination with polysorbate 60 as an emulsifier in mineral premixes and dietary supplements for animal feeds.

§ 573.980 Taurine.

The food additive taurine (2-aminoethanesulfonic acid) may be safely used in feed in accordance with the following prescribed conditions:

(a) It is used as a nutritional supplement in the feed of growing chickens.

(b) It is added to complete feeds so that the total taurine content does not exceed 0.054 percent of the feed.

(c) To assure safe use of the additive, the label and labeling shall bear in addition to the other information required by the Act:

(1) The name of the additive.

(2) The quantity of the additive contained therein.

(3) Adequate directions for use.

21 CFR Ch. I (4–1–11 Edition)

§ 573.1000 Verxite.

The food additive verxite may be safely used in animal feed in accordance with the following prescribed conditions:

(a) The additive is a magnesium-aluminum-iron silicate conforming to one of the following:

(1)(i) Verxite granules: The additive contains a minimum of 98 percent of hydrobiotite; it is thermally expanded and has a bulk density of from 5 to 9 pounds per cubic foot.

(ii) It is used or intended for use:

(a) In poultry feed at a level not to exceed 5 percent of the weight of the finished feed as a nonnutritive bulking agent for restricting calorie intake in pullet replacement feeds.

(b) As an anticaking or blending agent, pelleting aid, or nonnutritive carrier for the incorporation of nutrients in poultry, swine, dog, or ruminant feeds, in an amount not to exceed that necessary to accomplish its intended effect and in no case to exceed 1.5 percent of the dog feed or 5 percent of the final feed for other animals.

(2)(i) Verxite flakes: The additive contains a minimum of 98 percent of hydrobiotite; it has a bulk density of from 20 to 30 pounds per cubic foot.

(ii) It is used or intended for use as an anticaking or blending agent in ruminant feeds in an amount not to exceed that necessary to accomplish its intended effect and in no case to exceed 1 percent by weight of the final feed for ruminants.

(3)(i) Verxite grits: The additive contains a minimum of 80 percent of hydrobiotite; it has a bulk density of from 40 to 50 pounds per cubic foot.

(ii) It is used or intended for use as a partial roughage replacement in ruminant feeds in an amount not to exceed that necessary to accomplish its intended effect and in no case to exceed 1 percent by weight of the final feed.

(b) To assure safe use of the additive, the label of any feed additive supplement, feed additive concentrate, feed additive premix, or complete feed prepared therefrom shall bear, in addition to the other information required by the Act, the name of the additive (verxite granules, verxite flakes, or verxite grits), adequate directions for use, and, when the additive is present

Food and Drug Administration, HHS

§ 579.22

in excess of 1 percent, a statement of the quantity of the additive contained therein and the term “nonnutritive” in juxtaposition therewith.

§ 573.1010 Xanthan gum.

The food additive xanthan gum may be safely used in animal feed as follows:

(a) The food additive is xanthan gum as defined in §172.695 of this chapter and meets all of the specifications thereof.

(b) It is used or intended for use as a stabilizer, emulsifier, thickener, suspending agent, or bodying agent in animal feed as follows:

(1) In calf milk replacers at a maximum use level of 0.1 percent, as fed.

(2) In liquid feed supplements for ruminant animals at a maximum use level of 0.25 percent (5 pounds per ton).

(c) To assure safe use of the additive:

(1) The label of its container shall bear, in addition to other information required by the act, the name of the additive.

(2) The label or labeling of the additive container shall bear adequate directions for use.

[49 FR 44630, Nov. 8, 1984]

§ 573.1020 Yellow prussiate of soda.

Yellow prussiate of soda (sodium ferrocyanide decahydrate: $\text{Na}_4\text{Fe}(\text{Cn})_6 \cdot 10\text{H}_2\text{O}$) may be safely used as an anticaking agent in salt for animal consumption at a level not to exceed 13 parts per million. The additive contains a minimum of 99.0 percent by weight of sodium ferrocyanide decahydrate.

[41 FR 38657, Sept. 10, 1976; 41 FR 48100, Nov. 2, 1976]

PART 579—IRRADIATION IN THE PRODUCTION, PROCESSING, AND HANDLING OF ANIMAL FEED AND PET FOOD

Subpart A—General Provisions

Sec.

579.12 Incorporation of regulations in part 179.

Subpart B—Radiation and Radiation Sources

579.22 Ionizing radiation for treatment of animal diets.

579.40 Ionizing radiation for the treatment of poultry feed and poultry feed ingredients.

AUTHORITY: 21 U.S.C. 321, 342, 343, 348, 371.

Subpart A—General Provisions

§ 579.12 Incorporation of regulations in part 179.

Regulations providing for irradiation in the production, processing, and handling of food in part 179 of this chapter are incorporated in subchapter E as applicable to use in the production, processing, handling, and labeling of animal feed and pet food, except where specifically provided for in this part.

[51 FR 5993, Feb. 19, 1986]

Subpart B—Radiation and Radiation Sources

§ 579.22 Ionizing radiation for treatment of animal diets.

Ionizing radiation for treatment of complete diets for animals may be safely used under the following conditions:

(a) *Energy sources.* Ionizing radiation is limited to:

(1) Gamma rays for sealed units of the radionuclides cobalt-60 or cesium-137.

(2) Electrons generated from machine sources at energy levels not to exceed 10 million electron volts.

(b) *Uses.* (1) The ionizing radiation is used or intended for use in single treatment as follows:

Food for irradiation	Limitations	Use
Bagged complete diets, packaged feeds, feed ingredients, bulk feeds, animal treats and chews.	Absorbed dose: Not to exceed 50 kiloGrays. Feeds and feed ingredients treated by irradiation should be formulated to account for nutritional loss.	Microbial disinfection, control or elimination

(2) If an irradiated feed ingredient is less than 5 percent of the final product,