§ 180.421

Commodity	Parts per million
Hog, liver	0.1
Hog, meat	0.05
Hog, meat byproducts	0.05
Horse, fat	0.05
Horse, kidney	0.1
Horse, liver	0.1
Horse, meat	0.05
Horse, meat byproducts	0.05
Milk	0.05
Poultry, fat	0.05
Poultry, kidney	0.01
Poultry, liver	0.01
Poultry, meat	0.05
Poultry, meat byproducts	0.05
Sheep, fat	0.05
Sheep, kidney	0.1
Sheep, liver	0.1
Sheep, meat	0.05
Sheep, meat byproducts	0.05

- (b) Section 18 emergency exemptions. [Reserved]
- (c) Tolerances with regional registrations. [Reserved]
- (d) Indirect or inadvertent residues. Tolerances are established for indirect or inadvertent residues of the herbicide fluridone, including its metabolites and degradates, in or on the irrigated crop commodities and crop groupings in the table in this paragraph, resulting from use of irrigation water containing residues of 0.15 parts per million following applications of fluridone on or around aquatic sites. Where tolerances are established at higher levels from other uses of fluridone on the crops in the table in this paragraph, the higher tolerance also applies to residues in or on the irrigated commodity. Compliance with the tolerance levels specified in this paragraph is to be determined by measuring only fluridone, 1-methyl-3-phenyl-5-(3-

 $\label{eq:composition} \ensuremath{(\text{trifluoromethyl})\text{phenyl})-4(1H)-pyridinone, in or on the commodity.}$

Commodity	Parts per million
Animal feed, nongrass, group 18	0.15
Avocado	0.1
Berry, group 13	0.1
Cotton, undelinted seed	0.1
Cranberry	0.1
Fruit, citrus, group 10	0.1
Fruit, pome, group 11	0.1
Fruit, stone, group 12	0.1
Grain, cereal, forage, fodder and straw, group	
16	0.1
Grain, cereal, group 15	0.1
Grape	0.1
Grass, forage	0.15
Hop, dried cones	0.1
Nut, tree, group 14	0.1
Okra	0.1

Commodity	Parts per million
Strawberry	0.1
Vegetable, brassica, leafy, group 5	0.1
Vegetable, cucurbit, group 9	0.1
Vegetable, fruiting, group 8	0.1
Vegetable, leafy, except brassica, group 4	0.1
Vegetable, leaves of root and tuber, group 2	0.1
Vegetable, legume, group 6	0.1
Vegetable, root and tuber, group 1	0.1

[76 FR 23495, Apr. 27, 2011]

§ 180.421 Fenarimol; tolerances for residues.

(a) General. Tolerances are established for residues of fenarimol, including its metabolites and degradates, in or on the commodities in the following table. Compliance with the tolerance levels specified in the following table is to be determined by measuring only fenarimol alpha-(2 chlorophenyl)-alpha-(4-chlorophenyl)-5-pyrimidinemethanol.

Commodity	Parts per million
Apple	0.3
Apple, wet pomace	0.3
Banana	0.25
Cattle, fat	0.01
Cattle, kidney	0.01
Cattle, meat	0.01
Cattle, meat byproducts, except kidney	0.05
Cherry, sweet	1.0
Cherry, tart	1.0
Goat, fat	0.01
Goat, kidney	0.01
Goat, meat	0.01
Goat, meat byproducts, except kidney	0.05
Grape	0.1.
Hazelnut	0.02
Hop, dried cones	5.0
Horse, fat	0.01
Horse, kidney	0.01
Horse, meat	0.01
Horse, meat byproducts, except kidney	0.05
Pear	0.1
Pecan	0.02
Sheep, fat	0.01
Sheep, kidney	0.01
Sheep, meat	0.01
Sheep, meat byproducts, except kidney	0.05
Vegetable, cucurbit, group 9*	0.20

- *There are no U.S. registrations as of August 27, 2010.
- (b) Section 18 emergency exemptions. [Reserved]
- (c) Tolerances with regional registrations. [Reserved]

Environmental Protection Agency

(d) Indirect or inadvertent residues. [Reserved]

[51 FR 39662, Oct. 30, 1986, as amended at 53 FR 27349, July 20, 1988; 53 FR 44403, Nov. 3, 1988; 54 FR 45734, Oct. 31, 1989; 60 FR 33354, June 28, 1995; 62 FR 49937, Sept. 24, 1997; 62 FR 61447, Nov. 18, 1997; 67 FR 35048, May 17, 2002; 67 FR 41807, June 19, 2002; 69 FR 6567, Feb. 11, 2004; 71 FR 32846, June 7, 2006; 71 FR 54434, Sept. 15, 2006; 74 FR 68173, Dec. 23, 2009; 75 FR 56897, Sept. 17, 20101

§180.422 Tralomethrin; tolerances for residues.

(a) General. (1) Tolerances are established for the combined residues of the pesticide chemical tralomethrin (S)alpha-cyano-3-phenoxybenzyl (1R.3S)-2,2-dimethyl-3-[(RS)-1,2,2,2tetrabromoethyllcyclopropanecarboxylate) and its metabolites (S)-alpha-cyano-3phenoxybenzyl (1R,3R)-3(2,2dibromovinyl)-2,2-dimethylcyclopropanecarboxylate and (S)-alpha-

cyano-3-phenoxybenzyl(1S,3R)-3-(2,2dibromovinyl)-2,2-dimethyl-

cyclopropanecarboxylate calculated as the parent in or on the following food commodities:

Commodity	Parts per million
Broccoli	0.5
Cotton, undelinted seed	0.02
Cotton, oil	0.20
Lettuce, head	1.00
Lettuce, leaf	3.00
Soybean	0.05
Sunflower, seed	0.05

(2) A tolerance of 0.02 part per million is established for the combined insecticide residues of the tralomethrin ((S)-alpha-cyano-3phenoxybenzyl-(1R,3S)-2,2-dimethyl-3-[(RS)-1,2,2,2-tetrabromoethyl] cyclopropanecarboxylate) and its metabolites cis-deltamethrin [(S-alphacyano-3-phenoxybenzyl-(1R,3R)-3-[2,2dibromovinvl)-2.2dimethylcyclopropanecarboxylate] and trans-deltamethrin [(S)-alpha-cyano-3phenoxybenzyl dibromovinyl)-2,2-

(1S,3R)-3-(2,2dimethylcyclopropanecarboxylate] as

follows:

(i) In or on food commodities (other than those covered by a higher tolerance as a result of use on growing crops) in food-handling establishments.

- (ii) The insecticide may be present as residue from application tralomethrin in food-handling establishments, including food service, manufacturing, and processing establishments, such as restaurants, cafeterias, supermarkets, bakeries, breweries, dairies, meat slaughtering and packing plants, and canneries in accordance with the following prescribed condi-
- (A) Application shall be limited to a general surface and spot and/or crack and crevice treatment in food-handling establishments where food and food products are held, processed, prepared, and served. General surface application may be used only when the facility is not in operation provided exposed food has been covered or removed from the area being treated. All food-contact surfaces and equipment must be thoroughly cleaned after general surface applications. Spot and/or crack and crevice application may be used while the facility is in operation provided exposed food is covered or removed from the area being treated prior to application. Spray concentration shall be limited to a maximum of 0.06 percent active ingredient. Contamination of food and food-contact surfaces shall be avoided.
- (B) To assure safe use of the insecticide, its label and labelling shall conform to that registered with the U.S. Environmental Protection Agency and shall be used in accordance with such label and labelling.
- (3) A tolerance of 0.02 part per million is established for the combined insecticide residues of the ((S)-alpha-cyano-3tralomethrin phenoxybenzyl-(1R,3S)-2,2-dimethyl-3-[(RS)-1,2,2,2-tetrabromoethyl]cyclopropanecarboxylate) and its metabolites cis-deltamethrin [(S)-alphacyano-3-phenoxybenzyl-(1R,3R)-3-(2,2dibromovinyl)-2.2-

dimethylcyclopropanecarboxylate] and trans-deltamethrin [(S)-alpha-cyano-3phenoxybenzyl (1S,3R)-3-(2,2dibromovinyl)-2,2-

dimethylcyclopropanecarboxylate follows:

(i) In or on all feed items (other than those covered by a higher tolerance as a result of use on growing crops) in feed-handling establishments.