§ 180.460

§ 180.460 Benoxacor; tolerances for residues.

- (a) General. Tolerances are established for residues of the inert ingredient (safener) benoxacor (4-(dichloroacetyl)-3,4-dihydro-3-methyl-2H-1, 4-benzoxazine) at 0.01 parts per million (ppm) when used in pesticide formulations containing metolachlor or S-metolachlor in or on raw agricultural commodities for which tolerances have been established for metolachlor or S-metolachlor.
- (b) Section 18 energency exemptions. [Reserved]
- (c) Tolerances with regional registrations. [Reserved]
- (d) Indirect or inadvertent residues. [Reserved]

 $[63\ FR\ 7305,\ Feb.\ 13,\ 1998,\ as\ amended\ at\ 70\ FR\ 21631,\ Apr.\ 27,\ 2005]$

§ 180.461 Cadusafos; tolerances for residues.

A tolerance is established for the residues of the nematicide/insecticide cadusafos, *O*-ethyl *S,S*-di-*sec*-butyl phosphorodithioate, in or on the following raw agricultural commodity:

Commodity	Parts per million
Banana	0.01

There are no U.S. registrations as of May 10, 1994, for the nematicide/insecticid cadusafos.

[59 FR 39467, Aug. 3, 1994]

§ 180.462 Pyridate; tolerances for residues.

(a) General. Tolerances are established for residues of the herbicide pyridate, including its metabolites and degradates, in or on the commodities in the table in this paragraph. Compliance with the tolerance levels specified in this paragraph is to be determined by measuring only the sum of pyridate, O-(6-chloro-3-phenyl-4-pyridazinyl)-S-octyl-carbonothioate, and its metabolites, 6-chloro-3-phenyl-pyridazine-4-ol, calculated as the stoichiometric equivalent of pyridate, in or on the commodity.

Commodity	Parts per million
Brassica, head and stem, subgroup 5A	0.03
Cabbage	0.03
Chickpea, seed	0.1
Collards	0.03
Corn, field, forage	0.03
Corn, field, grain	0.03
Corn, field, stover	0.03
Corn, pop, grain	0.03
Corn, pop, stover	0.03
Peanut	0.03
Peppermint, tops	0.20
Spearmint, tops	0.20

- (b) Section 18 emergency exemptions. [Reserved]
- (c) Tolerances with regional registrations. [Reserved]
- (d) Indirect or inadvertent residues. [Reserved]

[57 FR 54303, Nov. 18, 1992, as amended at 62 FR 44558, Aug. 22, 1997; 63 FR 53844, Oct. 7, 1998; 64 FR 46298, Aug. 25, 1999; 65 FR 25652, May 3, 2000; 67 FR 35049, May 17, 2002; 72 FR 35665, June 29, 2007; 74 FR 46376, Sept. 9, 2009; 76 FR 23496, Apr. 27, 2011]

§ 180.463 Quinclorac; tolerances for residues.

(a) General. Tolerances are established for residues of the herbicide quinclorac, including its metabolites and degradates, in or on the commodities in the table in this paragraph. Compliance with the tolerance levels specified in this paragraph is to be determined by measuring only quinclorac, 3,7-dichloro-8-quinolinecarboxylic acid, in or on the commodity.

Commodity	Parts per million
Barley, grain	2.0
Cattle, fat	0.7
Cattle, meat byproducts	1.5
Cattle, meat	0.05
Egg	0.05
Goat, fat	0.7
Goat, meat byproducts	1.5
Goat, meat	0.05
Grain, aspirated fractions	1200
Grass, forage	150
Grass, hay	130
Hog, fat	0.7
Hog, meat byproducts	1.5
Hog, meat	0.05
Horse, fat	0.7
Horse, meat byproducts	1.5
Horse, meat	0.05
Milk	0.05
Poultry, fat	0.05
Poultry, meat byproducts	0.1
Poultry, meat	0.05
Rice, bran	15.0
Rice, grain	5.0