Environmental Protection Agency

§ 421.184 Standards of performance for new sources.

Any new source subject to this subpart shall achieve the following new source performance standards:

(a) Still liquor.

NSPS FOR THE PRIMARY AND SECONDARY GERMANIUM AND GALLIUM SUBCATEGORY

| Pollutant or pollutant property | Maximum for any 1 day | Maximum for monthly average |
|---------------------------------|--|-----------------------------------|
| | mg/kg (pounds per million pounds) of germanium chlorinated | |
| Arsenic | 131.70 | 58.59 |
| Lead | 26.46 | 12.60 |
| Zinc | 91.98 | 38.43 |
| Fluoride | 2,205.00 | 1,254.00 |
| Total suspended solids | 2,583.00 | 1,229.00 |
| pH | (1) | (1) |

¹ Within the range of 7.5 to 10.0 at all times.

(b) Chlorinator wet air pollution control.

NSPS FOR THE PRIMARY AND SECONDARY GERMANIUM AND GALLIUM SUBCATEGORY

| Pollutant or pollutant property | Maximum for any 1 day | Maximum for monthly average |
|---------------------------------|--|-----------------------------------|
| | mg/kg (pounds per million pounds) of germanium chlorinated | |
| Arsenic | 27.530 | 12.250 |
| Lead | 5.531 | 2.634 |
| Zinc | 19.230 | 8.034 |
| Fluoride | 461.000 | 262.100 |
| Total suspended solids | 540.000 | 256.800 |
| pH | (1) | (1) |

¹ Within the range of 7.5 to 10.0 at all times.

(c) Germanium hydrolysis filtrate.

NSPS FOR THE PRIMARY AND SECONDARY GERMANIUM AND GALLIUM SUBCATEGORY

| Pollutant or pollutant property | Maximum for 1 one day | Maximum for monthly average |
|--|--|---|
| | mg/kg pounds per millior pounds) of germanium hydrolyzed | |
| Arsenic Lead Zinc Fluoride Total suspended solids pH | 39.440 7.925 27.550 660.500 773.700 (¹) | 17.550 3.774 11.510 375.500 368.000 |

¹ Within the range of 7.5 to 10.0 at all times.

(d) Acid wash and rinse water.

NSPS FOR THE PRIMARY AND SECONDARY GERMANIUM AND GALLIUM SUBCATEGORY

| Pollutant or pollutant property | Maximum for any 1 day | Maximum for monthly average |
|---------------------------------|---|-----------------------------|
| | mg/kg (pounds per million pounds) of germanium washed | |
| Arsenic | 325.50 | 144.80 |
| Lead | 65.40 | 31.14 |
| Zinc | 227.40 | 94.99 |
| Fluoride | 5,450.00 | 3,099.00 |
| Total suspended solids | 6,385.00 | 3,037.00 |
| pH | (1) | (1) |

¹ Within the range of 7.5 to 10.0 at all times.

(e) Gallium hydrolysis filtrate.

NSPS FOR THE PRIMARY AND SECONDARY GERMANIUM AND GALLIUM SUBCATEGORY

| Pollutant or pollutant property | Maximum for any 1 day | Maximum for monthly average |
|---------------------------------|---|-----------------------------------|
| | mg/kg (pounds per million pounds) of gallium hydrolyzed | |
| Arsenic | 70.450 | 31.350 |
| Lead | 14.160 | 6.742 |
| Zinc | 49.220 | 20.560 |
| Fluoride | 1,180.000 | 670.800 |
| Total suspended solids | 1,382.000 | 657.300 |
| pH | (1) | (1) |

¹ Within the range of 7.5 to 10.0 at all times.

(f) Solvent extraction raffinate.

NSPS FOR THE PRIMARY AND SECONDARY GERMANIUM AND GALLIUM SUBCATEGORY

| Pollutant or pollutant property | Maximum for any 1 day | Maximum for monthly average |
|---------------------------------|--|---|
| | mg/kg (pounds per millior pounds) of gallium pro duced by solvent extrac tion | |
| Arsenic | 39.330 7.904 27.480 658.700 771.600 | 17.500 3.764 11.480 374.500 367.000 |
| pH | (1) | (1) |

¹ Within the range of 7.5 to 10.0 at all times.

§421.185 Pretreatment standards for existing sources.

Except as provided in 40 CFR 403.7 and 403.13, any existing source subject to this subpart which introduces pollutants into a publicly owned treatment works must comply with 40 CFR part 403 and achieve the following pretreatment standards for existing