§ 493.1264

(commercially prepared), and shipment of lactophenol cotton blue when prepared or opened for intended reactivity with a control organism(s).

- (b) For antifungal susceptibility tests, the laboratory must check each batch of media and each lot number and shipment of antifungal agent(s) before, or concurrent with, initial use, using an appropriate control organism(s).
- (1) The laboratory must establish limits for acceptable control results.
- (2) Each day tests are performed, the laboratory must use the appropriate control organism(s) to check the procedure
- (3) The results for the control organism(s) must be within established limits before reporting patient results.
- (c) The laboratory must document all control procedures performed, as specified in this section.

§ 493.1264 Standard: Parasitology.

- (a) The laboratory must have available a reference collection of slides or photographs and, if available, gross specimens for identification of parasites and use these references in the laboratory for appropriate comparison with diagnostic specimens.
- (b) The laboratory must calibrate and use the calibrated ocular micrometer for determining the size of ova and parasites, if size is a critical parameter.
- (c) Each month of use, the laboratory must check permanent stains using a fecal sample control material that will demonstrate staining characteristics.
- (d) The laboratory must document all control procedures performed, as specified in this section.

§ 493.1265 Standard: Virology.

- (a) When using cell culture to isolate or identify viruses, the laboratory must simultaneously incubate a cell substrate control or uninoculated cells as a negative control material.
- (b) The laboratory must document all control procedures performed, as specified in this section.

§ 493.1267 Standard: Routine chemistry.

For blood gas analyses, the laboratory must perform the following:

- (a) Calibrate or verify calibration according to the manufacturer's specifications and with at least the frequency recommended by the manufacturer.
- (b) Test one sample of control material each 8 hours of testing using a combination of control materials that include both low and high values on each day of testing.
- (c) Test one sample of control material each time specimens are tested unless automated instrumentation internally verifies calibration at least every 30 minutes.
- (d) Document all control procedures performed, as specified in this section.

§ 493.1269 Standard: Hematology.

- (a) For manual cell counts performed using a hemocytometer—
- (1) One control material must be tested each 8 hours of operation; and
- (2) Patient specimens and control materials must be tested in duplicate.
- (b) For all nonmanual coagulation test systems, the laboratory must include two levels of control material each 8 hours of operation and each time a reagent is changed.
 - (c) For manual coagulation tests—
- (1) Each individual performing tests must test two levels of control materials before testing patient samples and each time a reagent is changed; and
- (2) Patient specimens and control materials must be tested in duplicate.
- (d) The laboratory must document all control procedures performed, as specified in this section.

§ 493.1271 Standard: Immunohematology.

- (a) Patient testing. (1) The laboratory must perform ABO grouping, D(Rho) typing, unexpected antibody detection, antibody identification, and compatibility testing by following the manufacturer's instructions, if provided, and as applicable, 21 CFR 606.151(a) through (e).
- (2) The laboratory must determine ABO group by concurrently testing unknown red cells with, at a minimum, anti-A and anti-B grouping reagents. For confirmation of ABO group, the unknown serum must be tested with known A1 and B red cells.

- (3) The laboratory must determine the D(Rho) type by testing unknown red cells with anti-D (anti-Rho) blood typing reagent.
- (b) Immunohematological testing and distribution of blood and blood products. Blood and blood product testing and distribution must comply with 21 CFR 606.100(b)(12); 606.160(b)(3)(ii) and (b)(3)(v); 610.40; 640.5(a), (b), (c), and (e); and 640.11(b).
- (c) Blood and blood products storage. Blood and blood products must be stored under appropriate conditions that include an adequate temperature alarm system that is regularly inspected.
- (1) An audible alarm system must monitor proper blood and blood product storage temperature over a 24-hour period.
- (2) Inspections of the alarm system must be documented.
- (d) Retention of samples of transfused blood. According to the laboratory's established procedures, samples of each unit of transfused blood must be retained for further testing in the event of transfusion reactions. The laboratory must promptly dispose of blood not retained for further testing that has passed its expiration date.
- (e) Investigation of transfusion reactions. (1) According to its established procedures, the laboratory that performs compatibility testing, or issues blood or blood products, must promptly investigate all transfusion reactions occurring in facilities for which it has investigational responsibility and make recommendations to the medical staff regarding improvements in transfusion procedures.
- (2) The laboratory must document, as applicable, that all necessary remedial actions are taken to prevent recurrences of transfusion reactions and that all policies and procedures are reviewed to assure they are adequate to ensure the safety of individuals being transfused.
- (f) Documentation. The laboratory must document all control procedures performed, as specified in this section.

[68 FR 3703, Jan. 24, 2003; 68 FR 50724, Aug. 22, 2003]

§ 493.1273 Standard: Histopathology.

- (a) As specified in §493.1256(e)(3), fluorescent and immunohistochemical stains must be checked for positive and negative reactivity each time of use. For all other differential or special stains, a control slide of known reactivity must be stained with each patient slide or group of patient slides. Reaction(s) of the control slide with each special stain must be documented.
- (b) The laboratory must retain stained slides, specimen blocks, and tissue remnants as specified in § 493.1105. The remnants of tissue specimens must be maintained in a manner that ensures proper preservation of the tissue specimens until the portions submitted for microscopic examination have been examined and a diagnosis made by an individual qualified under §§ 493.1449(b), (l), or (m).
- (c) An individual who has successfully completed a training program in neuromuscular pathology approved by HHS may examine and provide reports for neuromuscular pathology.
- (d) Tissue pathology reports must be signed by an individual qualified as specified in paragraph (b) or, as appropriate, paragraph (c) of this section. If a computer report is generated with an electronic signature, it must be authorized by the individual who performed the examination and made the diagnosis.
- (e) The laboratory must use acceptable terminology of a recognized system of disease nomenclature in reporting results.
- (f) The laboratory must document all control procedures performed, as specified in this section.

 $[68~{\rm FR}~3703,~{\rm Jan.}~24,~2003;~68~{\rm FR}~50724,~{\rm Aug.}~22,~2003]$

§ 493.1274 Standard: Cytology.

- (a) Cytology slide examination site. All cytology slide preparations must be evaluated on the premises of a laboratory certified to conduct testing in the subspecialty of cytology.
- (b) Staining. The laboratory must have available and follow written policies and procedures for each of the following, if applicable:
- (1) All gynecologic slide preparations must be stained using a Papanicolaou