

shape of kernels of rice that are broken as prescribed in FGIS instructions.

(n) *Ungelatinized kernels.* Whole or large broken kernels of parboiled rice with distinct white or chalky areas due to incomplete gelatinization of the starch.

(o) *Whole and large broken kernels.* Rice (including seeds) that (1) passes over a 6 plate (for southern production), or (2) remains on top of a 6 sieve (for western production).

(p) *Whole kernels.* Unbroken kernels of rice and broken kernels of rice which are at least three-fourths of an unbroken kernel.

(q) *6 sieve.* A metal sieve 0.032-inch thick, perforated with rows of round holes 0.0938 ( $\frac{3}{64}$ ) inch in diameter.

(r) *6 plate.* A laminated metal plate 0.142-inch thick, with a top lamina 0.051-inch thick, perforated with rows of round holes 0.0938 ( $\frac{3}{64}$ ) inch in diameter, and a bottom lamina 0.091-inch thick, without perforations.

[42 FR 40869, Aug. 12, 1977; 42 FR 64356, Dec. 23, 1977, as amended at 47 FR 34516, Aug. 10, 1982; 54 FR 21403, May 18, 1989; 54 FR 51344, Dec. 14, 1989. Redesignated at 60 FR 16364, Mar. 30, 1995]

#### PRINCIPLES GOVERNING APPLICATION OF STANDARDS

##### § 868.203 Basis of determination.

The determination of seeds, objectionable seeds, heat-damaged kernels, red rice and damaged kernels, chalky kernels, other types, color, and the special grade Parboiled rough rice shall be on the basis of the whole and large broken kernels of milled rice that are produced in the milling of rough rice to a well-milled degree. When determining class, the percentage of (a) whole kernels of rough rice shall be determined on the basis of the original sample, and (b) types of rice shall be determined on the basis of the whole and large broken kernels of milled rice that are produced in the milling of rough rice to a well-milled degree. Smutty kernels shall be determined on the basis of the rough rice after it has been cleaned and shelled as prescribed in FGIS instructions, or by any method that is approved by the Administrator as giving equivalent results. All other determinations shall be on the basis of the

original sample. Mechanical sizing of kernels shall be adjusted by handpicking as prescribed in FGIS instructions, or by any method that is approved by the Administrator as giving equivalent results.

[42 FR 40869, Aug. 12, 1977; 42 FR 64356, Dec. 23, 1977, as amended at 47 FR 34516, Aug. 10, 1982; 54 FR 21403, May 18, 1989. Redesignated at 60 FR 16364, Mar. 30, 1995]

##### § 868.204 Interpretive line samples.

Interpretive line samples showing the official scoring line for factors that are determined by visual examinations shall be maintained by the Federal Grain Inspection Service, U.S. Department of Agriculture, and shall be available for reference in all inspection offices that inspect and grade rice.

[42 FR 40869, Aug. 12, 1977; 42 FR 64356, Dec. 23, 1977, as amended at 47 FR 34516, Aug. 10, 1982. Redesignated at 54 FR 21403, May 18, 1989, and 60 FR 16364, Mar. 30, 1995]

##### § 868.205 Milling requirements.

In determining milling yield (see § 868.202(f)) in rough rice, the degree of milling shall be equal to, or better than, that of the interpretive line sample for "well-milled" rice.

[42 FR 40869, Aug. 12, 1977. Redesignated at 54 FR 21413, May 18, 1989, and further redesignated and amended at 60 FR 16364, 16365, Mar. 30, 1995]

##### § 868.206 Milling yield determination.

Milling yield shall be determined by the use of an approved device in accordance with procedures prescribed in FGIS instructions. For the purpose of this paragraph, "approved device" shall include the McGill Miller No. 3 and any other equipment that is approved by the Administrator as giving equivalent results.

NOTE: Milling yield shall not be determined when the moisture content of the rough rice exceeds 18.0 percent.

[42 FR 40869, Aug. 12, 1977; 42 FR 64356, Dec. 23, 1977, as amended at 47 FR 34516, Aug. 10, 1982; Redesignated and amended at 54 FR 21403, May 18, 1989, and further redesignated at 60 FR 16364, Mar. 30, 1995]

##### § 868.207 Moisture.

Water content in rough rice as determined by an approved device in accordance with procedures prescribed in the