21 CFR Ch. I (4–1–12 Edition)

until hatch for all warmwater species of freshwater-reared finfish eggs.

(ii) Freshwater-reared salmonids: 100 mg/L for 30 minutes or 50 to 100 mg/L for 60 minutes once per day on alternate days for three treatments in a continuous flow water supply or as a static bath.

(iii) Coolwater species of freshwaterreared finfish fingerlings and adults (except northern pike & paddlefish) and channel catfish fingerlings and adults: 50 to 75 mg/L for 60 minutes once per day on alternate days for three treatments in continuous flow water supply or as a static bath. Coolwater species of freshwater-reared finfish fry (except northern pike, pallid sturgeon & paddlefish) and channel catfish fry: 50 mg/L for 60 minutes once per day on alternate days for three treatments in continuous flow water supply or as a static bath.

(2) Indications for use. For control of mortality in freshwater-reared finfish eggs due to saprolegniasis; for control of mortality in freshwater-reared salmonids due to bacterial gill disease associated with Flavobacterium branchiophilum; and for control of mortality in freshwater-reared coolwater finfish and channel catfish due to external columnaris disease associated Flavobacteriumcolumnarewith (Flexibacter columnaris).

(3) Limitations. Initial bioassay on a small number is recommended before treating the entire group. Eggs: Some strains of rainbow trout eggs are sensitive to hydrogen peroxide treatment at a time during incubation concurrent with blastopore formation through closure, about 70 to 140 Daily Temperature Units, °C. Consider withholding treatalternate ment or using an therapeutant during that sensitive time to reduce egg mortalities due to drug toxicity. Finfish: Use with caution on walleye. Preharvest withdrawal time: zero days.

[72 FR 5330, Feb. 6, 2007]

## §529.1186 Isoflurane.

(a) *Specifications*. The drug is a clear, colorless, stable liquid.

(b) *Sponsors*. See Nos. 000074, 000209, 010019, 012164, 065085, and 066794 in §510.600(c) of this chapter.

(c) *Conditions of use*. Administer by inhalation:

(1) Amount—(i) Horses: For induction of surgical anesthesia: 3 to 5 percent isoflurane (with oxygen) for 5 to 10 minutes. For maintenance of surgical anesthesia: 1.5 to 1.8 percent isoflurane (with oxygen).

(ii) *Dogs:* For induction of surgical anesthesia: 2 to 2.5 percent isoflurane (with oxygen) for 5 to 10 minutes. For maintenance of surgical anesthesia: 1.5 to 1.8 percent isoflurane (with oxygen).

(2) *Indications for use*. For induction and maintenance of general anesthesia in horses and dogs.

(3) *Limitations*. Do not use in horses intended for human consumption. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[51 FR 594, Jan. 7, 1986, as amended at 54 FR 23472, June 1, 1989; 58 FR 17346, Apr. 2, 1993; 59 FR 44315, Aug. 29, 1994; 60 FR 40456, Aug. 9, 1995; 63 FR 8122, Feb. 18, 1998; 63 FR 24106, May 1, 1998; 66 FR 17510, Apr. 2, 2001; 71 FR 43967, Aug. 3, 2006; 74 FR 68530, Dec. 28, 2009; 76 FR 16533, Mar. 24, 2011]

## § 529.1660 Oxytetracycline.

(a) *Specifications*—(1) Each gram of powder contains 366 milligrams (mg) oxytetracycline hydrochloride.

(2) Each gram of powder contains 753 mg oxytetracycline hydrochloride.

(b) *Sponsors.* See sponsors in §510.600(c) of this chapter for use of products described in paragraph (a) of this section as in paragraph (d) of this section.

(1) Nos. 046573 and 061623 for use of product in paragraph (a)(1) of this section.

(2) Nos. 000069, 048164, and 059130 for use of product described in paragraph (a)(2) of this section.

(c) Related tolerances. See \$556.500 of this chapter.

(d) Conditions of use in finfish—(1) Amount. Immerse fish in a solution containing 200 to 700 mg oxytetracycline hydrochloride (buffered) per liter of water for 2 to 6 hours.

(2) *Indications for use*. For skeletal marking of finfish fry and fingerlings.

[69 FR 6557, Feb. 11, 2004, as amended at 69
FR 61999, Oct. 22, 2004; 70 FR 41140, July 18, 2005; 72 FR 26289, May 9, 2007; 76 FR 17026, Mar. 28, 2011]