

an amount not to exceed 1.0 percent by weight of the finished feed.

(d) *Labeling.* The labels and labeling shall bear, in addition to the other information required by the Act:

(1) The name of the additive, hydrolyzed leather meal.

(2) Adequate directions to provide finished feeds complying with paragraph (c) of this section.

**§ 573.560 Iron ammonium citrate.**

Iron ammonium citrate may be safely used in animal feed in accordance with the following prescribed conditions:

(a) The additive is the chemical green ferric ammonium citrate.

(b) The additive is used or intended for use as an anticaking agent in salt for animal consumption so that the level of iron ammonium citrate does not exceed 25 parts per million (0.0025 percent) in the finished salt.

(c) To assure safe use of the additive the label or labeling of the additive shall bear, in addition to the other information required by the Act:

(1) The name of the additive.

(2) Adequate directions to provide a final product that complies with the limitations prescribed in paragraph (b) of this section.

**§ 573.580 Iron-choline citrate complex.**

Iron-choline citrate complex made by reacting approximately equimolecular quantities of ferric hydroxide, choline, and citric acid may be safely used as a source of iron in animal feed.

**§ 573.600 Lignin sulfonates.**

Lignin sulfonates may be safely used in animal feeds in accordance with the following prescribed conditions:

(a) For the purpose of this section, the food additive is either one, or a combination of, the ammonium, calcium, magnesium, or sodium salts of the extract of spent sulfite liquor derived from the sulfite digestion of wood or of abaca (*Musa textilis*) or of sisal (*Agave sisalana*) in either a liquid form (moisture not to exceed 50 percent by weight) or dry form (moisture not to exceed 6 percent by weight).

(b) It is used or intended for use in an amount calculated on a dry weight basis, as follows:

(1) As a pelleting aid in the liquid or dry form in an amount not to exceed 4 percent of the finished pellets.

(2) As a binding aid in the liquid form in the flaking of feed grains in an amount not to exceed 4 percent of the flaked grain.

(3) As a surfactant in molasses used in feeds, as liquid lignin sulfonate, in an amount not to exceed 11 percent of the molasses.

(4) As a source of metabolizable energy, in the liquid or dry form, in an amount not to exceed 4 percent of the finished feed.

**§ 573.620 Menadione dimethylpyrimidinol bisulfite.**

The food additive, menadione dimethylpyrimidinol bisulfite, may be safely used in accordance with the following conditions:

(a) The additive is the 2-hydroxy-4,6-dimethylpyrimidinol salt of menadione (C<sub>17</sub>H<sub>18</sub>O<sub>6</sub>N<sub>2</sub>S).

(b) The additive is used or intended for use as a nutritional supplement for the prevention of vitamin K deficiency as follows:

(1) In chicken and turkey feed at a level not to exceed 2 grams per ton of complete feed.

(2) In the feed of growing and finishing swine at a level not to exceed 10 grams per ton of feed.

(c) To assure safe use, the label and labeling of the additive shall bear adequate directions for use.

**§ 573.625 Menadione nicotinamide bisulfite.**

The food additive may be safely used as follows:

(a) The additive is 1,2,3,4-tetrahydro-2-methyl-1,4-dioxo-2-naphthalene sulfonic acid with 3-pyridine carboxylic acid amine (CAS No. 73581-79-0).

(b) The additive is used or intended for use as a nutritional supplement for both the prevention of vitamin K deficiency and as a source of supplemental niacin as follows:

(1) In chicken and turkey feeds at a level not to exceed 2 grams per ton of complete feed.

(2) In growing and finishing swine feeds at a level not to exceed 10 grams per ton of complete feed.