

§ 640.64

21 CFR Ch. I (4-1-12 Edition)

donor under the provisions of this section;

(3) A blood hemoglobin level of no less than 12.5 grams of hemoglobin per 100 milliliters of blood or a hematocrit level of 38 percent;

(4) A normal pulse rate;

(5) A total serum or total plasma protein of no less than 6.0 grams per 100 milliliters of blood;

(6) Weight, which shall be at least 110 pounds;

(7) Freedom from acute respiratory diseases;

(8) Freedom from any infectious skin disease at the site of phlebotomy and from any such disease generalized to such an extent as to create a risk of contamination of the plasma;

(9) Freedom from any disease, other than malaria, transmissible by blood transfusion, insofar as can be determined by history and examinations indicated in this section;

(10) Freedom of the arms and forearms from skin punctures or scars indicative of addiction to self-injected narcotics;

(11) Freedom from a history of viral hepatitis after the 11th birthday;

(12) Freedom from a history of close contact within 12 months of donation with an individual having viral hepatitis;

(13) Freedom from a history of having received, within 12 months, human blood or any derivative of human blood which the Food and Drug Administration has advised the blood establishment is a possible source of viral hepatitis, except for specific immunization performed in accordance with § 640.66.

(d) *General.* Any donor who, in the opinion of the interviewer, appears to be under the influence of any drug, alcohol, or for any reason does not appear to be providing reliable answers to medical history questions, shall not be considered a suitable donor.

(e) *Failure to return red blood cells.* Any donor who has not had the red blood cells returned from a unit of blood collected during a plasmapheresis procedure or who has been a donor of a unit of whole blood shall not be subjected to plasmapheresis for a period of 8 weeks, unless:

(1) The donor has been examined by a qualified licensed physician and cer-

tified by the physician to be acceptable for further plasmapheresis before expiration of the 8-week period;

(2) The donor possesses an antibody that is (i) transitory, (ii) of a highly unusual or infrequent specificity, or (iii) of an unusually high titer; and

(3) The special characteristics of the antibody and the need for plasmapheresing the donor are documented.

[38 FR 32089, Nov. 20, 1973, as amended at 41 FR 10768, Mar. 12, 1976; 43 FR 9805, Mar. 10, 1978; 43 FR 12311, Mar. 24, 1978; 46 FR 57480, Nov. 24, 1981; 50 FR 4140, Jan. 29, 1985; 64 FR 45373, Aug. 19, 1999; 66 FR 1837, Jan. 10, 2001; 66 FR 40890, Aug. 6, 2001]

§ 640.64 Collection of blood for Source Plasma.

(a) *Supervision.* All blood for the collection of Source Plasma shall be drawn from the donor by a qualified licensed physician or by persons under his supervision trained in the procedure.

(b) *Blood containers.* Blood containers and donor sets must be pyrogen-free, sterile, and identified by lot number.

(c) *The anticoagulant solution.* The anticoagulant solution must be sterile and pyrogen-free. Anticoagulant solutions must be compounded and used according to a formula that has been approved for the applicant by the Director, Center for Biologics Evaluation and Research.

(d) *Donor identification.* Each unit of blood and plasma shall be so marked or identified by number or other symbol so as to relate it directly to the donor.

(e) *Prevention of contamination of the blood and plasma.* The skin of the donor at the site of phlebotomy shall be prepared thoroughly and carefully by a method that gives maximum assurance of a sterile container of blood. The blood shall be collected, the plasma separated, and the cells returned to the donor by aseptic methods in a sterile system which may be closed, or may be vented if the vent protects the blood

cells and plasma against contamination.

[38 FR 32089, Nov. 20, 1973; 39 FR 13632, Apr. 16, 1974, as amended at 41 FR 10768, Mar. 12, 1976; 49 FR 23834, June 8, 1984; 50 FR 4140, Jan. 29, 1985; 55 FR 11013, Mar. 26, 1990; 59 FR 49351, Sept. 28, 1994; 63 FR 16685, Apr. 6, 1998; 64 FR 56453, Oct. 20, 1999; 72 FR 45888, Aug. 16, 2007]

§ 640.65 Plasmapheresis.

(a) *Procedure-general.* The plasmapheresis procedure is a procedure in which, during a single visit to the establishment, blood is removed from a donor, the plasma separated from the formed elements, and at least the red blood cells returned to the donor. This procedure shall be described in detail in the biologics license application.

(b) *Procedures-specific requirements.* The plasmapheresis procedure shall meet the following requirements:

(1)(i) A sample of blood shall be drawn from each donor on the day of the first medical examination or plasmapheresis, whichever comes first and at least every 4 months thereafter by a qualified licensed physician or by persons under his supervision and trained in such procedure. A serologic test for syphilis, a total plasma or serum protein determination, and a plasma or serum protein electrophoresis or quantitative immuno-diffusion test or an equivalent test to determine immunoglobulin composition of the plasma or serum shall be performed on the sample.

(ii) A repeat donor who does not return for plasmapheresis at the time the 4-month sample is due to be collected may be plasmapheresed on the day he appears: *Provided*, That no longer than 6 months has elapsed since the last sample was collected, and the physician on the premises approves the plasmapheresis procedure and so indicates by signing the donor's record before such procedure is performed. The sample for the 4-month tests shall be collected on the day of the donor's return.

(iii) A repeat donor from whom the plasmapheresis center is unable to obtain a sample for testing as prescribed in paragraph (b)(1)(i) of this section for a total period exceeding 6 months shall be processed as a new donor.

(2)(i) The accumulated laboratory data, including tracings, if any, of the plasma or serum protein electrophoresis pattern, the calculated values of each component, and the collection records shall be reviewed by a qualified licensed physician within 21 days after the sample is drawn to determine whether or not the donor may continue in the program. The review shall be signed by the reviewing physician. If the protein composition is not within normal limits established by the testing laboratory, or if the total protein is less than 6.0 grams per 100 milliliters of samples, the donor shall be removed from the program until these values return to normal.

(ii) A donor with a reactive serologic test for syphilis shall not be plasmapheresed again until the donor's serum is tested and found to be non-reactive to a serologic test for syphilis, except as provided in paragraph (b)(2)(iii) and (iv) of this section.

(iii) A donor whose serum is determined to have a biologic false-positive reaction to a serologic test for syphilis may be plasmapheresed: *Provided*, That the donor's file identifies the serologic test for syphilis and results used to confirm the biologic false-positive reaction and indicates that the physician on the premises has determined the false-positive reaction is not the result of an underlying disorder that would disqualify the donor from participation in the plasmapheresis program. If the serologic test for syphilis is performed at a facility other than the plasmapheresis center, all applicable provisions of § 640.71 shall be met.

(iv) A donor with a reactive serologic test for syphilis may be plasmapheresed only to obtain plasma to be used for further manufacturing into control serum for the serologic test for syphilis: *Provided*, That the physician on the premises approves the donation, the donor's file contains a signed statement from a physician or clinic establishing that treatment for syphilis has been initiated and that continuance in the plasmapheresis program will not interfere with or jeopardize the treatment of the syphilitic donor.

(3) A donor identification system shall be established that positively