

**§ 352.77**

(3) 20-minute rest period (do not towel test sites).

(4) 20 minutes moderate activity in water.

(5) Conclude water test (air dry test sites without toweling).

(6) Begin solar simulator exposure to test site areas as described in §352.73.

(b) *Procedure for testing a very water resistant sunscreen product.* For sunscreen products making the claim of "very water resistant," the label SPF shall be the label SPF value determined after 80 minutes of water immersion using the following procedure for the very water resistant test:

(1) Apply sunscreen product (followed by the waiting period after application of the sunscreen product indicated on the product labeling).

(2) 20 minutes moderate activity in water.

(3) 20-minute rest period (do not towel test sites).

(4) 20 minutes moderate activity in water.

(5) 20-minute rest period (do not towel test sites).

(6) 20 minutes moderate activity in water.

(7) 20-minute rest period (do not towel test sites).

(8) 20 minutes moderate activity in water.

(9) Conclude water test (air dry test sites without toweling).

(10) Begin solar simulator exposure to test site areas as described in §352.73.

**§ 352.77 Test modifications.**

The formulation or mode of administration of certain products may require modification of the testing procedures in this subpart. In addition, alternative methods (including automated or in vitro procedures) employing the same basic procedures as those described in this subpart may be used. Any proposed modification or alternative procedure shall be submitted as a petition in accord with §10.30 of this chapter. The petition should contain data to support the modification or data demonstrating that an alternative procedure provides results of equivalent accuracy. All information submitted will be subject to the disclosure rules in part 20 of this chapter.

**21 CFR Ch. I (4-1-13 Edition)**

**PART 355—ANTICARIES DRUG PRODUCTS FOR OVER-THE-COUNTER HUMAN USE**

**Subpart A—General Provisions**

- Sec.
- 355.1 Scope.
- 355.3 Definitions.

**Subpart B—Active Ingredients**

- 355.10 Anticaries active ingredients.
- 355.20 Packaging conditions.

**Subpart C—Labeling**

- 355.50 Labeling of anticaries drug products.
- 355.55 Principal display panel of all fluoride rinse drug products.
- 335.60 Professional labeling.

**Subpart D—Testing Procedures**

- 355.70 Testing procedures for fluoride dentifrice drug products.

AUTHORITY: 21 U.S.C. 321, 351, 352, 353, 355, 360, 371.

SOURCE: 60 FR 52507, Oct. 6, 1995, unless otherwise noted.

EDITORIAL NOTE: Nomenclature changes to part 355 appear at 69 FR 13717, Mar. 24, 2004.

**Subpart A—General Provisions**

**§ 355.1 Scope.**

(a) An over-the-counter anticaries drug product in a form suitable for topical administration to the teeth is generally recognized as safe and effective and is not misbranded if it meets each condition in this part and each general condition established in §330.1 of this chapter.

(b) References in this part to regulatory sections of the Code of Federal Regulations are to chapter I of title 21 unless otherwise noted.

**§ 355.3 Definitions.**

As used in this part:

(a) *Abrasive.* Solid materials that are added to dentifrices to facilitate mechanical removal of dental plaque, debris, and stain from tooth surfaces.

(b) *Anhydrous glycerin.* An ingredient that may be prepared by heating glycerin U.S.P. at 150 °C for 2 hours to drive off the moisture content.

(c) *Anticaries drug.* A drug that aids in the prevention and prophylactic

treatment of dental cavities (decay, caries).

(d) *Dental caries*. A disease of calcified tissues of teeth characterized by demineralization of the inorganic portion and destruction of the organic matrix.

(e) *Dentifrice*. An abrasive-containing dosage form (gel, paste, or powder) for delivering an anticaries drug to the teeth.

(f) *Fluoride*. The inorganic form of the chemical element fluorine in combination with other elements.

(g) *Fluoride ion*. The negatively charged atom of the chemical element fluorine.

(h) *Fluoride supplement*. A special treatment rinse dosage form that is intended to be swallowed, and is promoted to health professionals for use in areas where the water supply contains 0 to 0.7 parts per million (ppm) fluoride ion.

(i) *Preventive treatment gel*. A dosage form for delivering an anticaries drug to the teeth. Preventive treatment gels are formulated in an anhydrous glycerin base with suitable thickening agents included to adjust viscosity. Preventive treatment gels do not contain abrasives.

(j) *Treatment rinse*. A liquid dosage form for delivering an anticaries drug to the teeth.

(k) *Treatment rinse concentrated solution*. A fluoride treatment rinse in a concentrated form to be mixed with water before using to result in the appropriate fluoride concentration specified in the monograph.

(l) *Treatment rinse effervescent tablets*. A fluoride treatment rinse prepared by adding an effervescent tablet (a concentrated solid dosage form) to water before using to result in the appropriate fluoride concentration specified in the monograph.

(m) *Treatment rinse powder*. A fluoride treatment rinse prepared by adding the powder (a concentrated solid dosage form) to water before using to result in the appropriate fluoride concentration specified in the monograph.

[60 FR 52507, Oct. 6, 1995, as amended at 61 FR 52286, Oct. 7, 1996]

## Subpart B—Active Ingredients

### § 355.10 Anticaries active ingredients.

The active ingredient of the product consists of any of the following when used in the concentration and dosage form established for each ingredient:

(a) *Sodium fluoride*—(1) *Dentifrices containing 850 to 1,150 ppm theoretical total fluorine in a gel or paste dosage form*. Sodium fluoride 0.188 to 0.254 percent with an available fluoride ion concentration  $\geq$  650 parts per million (ppm).

(2) *Dentifrices containing 850 to 1,150 ppm theoretical total fluorine in a powdered dosage form*. Sodium fluoride 0.188 to 0.254 percent with an available fluoride ion concentration of  $\geq$  850 ppm for products containing the abrasive sodium bicarbonate and a poured-bulk density of 1.0 to 1.2 grams per milliliter.

(3) *Treatment rinses*. (i) An aqueous solution of acidulated phosphate fluoride derived from sodium fluoride acidulated with a mixture of sodium phosphate, monobasic, and phosphoric acid to a level of 0.1 molar phosphate ion and a pH of 3.0 to 4.5 and which yields an effective fluoride ion concentration of 0.02 percent.

(ii) An aqueous solution of acidulated phosphate fluoride derived from sodium fluoride acidulated with a mixture of sodium phosphate, dibasic, and phosphoric acid to a pH of 3.5 and which yields an effective fluoride ion concentration of 0.01 percent.

(iii) Sodium fluoride 0.02 percent aqueous solution with a pH of approximately 7.

(iv) Sodium fluoride 0.05 percent aqueous solution with a pH of approximately 7.

(v) Sodium fluoride concentrate containing adequate directions for mixing with water before using to result in a 0.02-percent or 0.05-percent aqueous solution with a pH of approximately 7.

(b) *Sodium monofluorophosphate*—(1) *Dentifrices containing 850 to 1,150 ppm theoretical total fluorine in a gel or paste dosage form*. Sodium monofluorophosphate 0.654 to 0.884 percent with an available fluoride ion concentration (consisting of  $\text{PO}_3\text{F}^-$  and  $\text{F}^-$  combined)  $\geq$  800 ppm.

(2) *Dentifrices containing 1,500 ppm theoretical total fluorine in a gel or paste*