

producing eggs for human consumption. Do not administer within 5 days of slaughter.

(3) *Swine*—(i) *Amount*. 250 mg per gallon as the only source of drinking water for 3 to 10 days, depending on the severity of the condition being treated.

(ii) *Indications for use*. (A) For the treatment and control of swine dysentery associated with *Brachyspira hyodysenteriae* when followed immediately by tylosin phosphate medicated feed; and for the control of porcine proliferative enteropathies (PPE, ileitis) associated with *Lawsonia intracellularis* when followed immediately by tylosin phosphate medicated feed.

(B) For the treatment and control of swine dysentery associated with *Brachyspira hyodysenteriae*.

(iii) *Limitations*. Prepare a fresh solution daily. Do not administer within 48 hours of slaughter. As indicated in paragraph (d)(3)(ii)(A) of this section, follow with tylosin phosphate medicated feed as in §558.625(f)(1)(vi)(c) of this chapter.

(4) *Honey bees*—(i) *Amount*. Mix 200 milligrams tylosin in 20 grams confecttioners' powdered sugar. Use immediately. Apply (dust) this mixture over the top bars of the brood chamber once weekly for 3 weeks.

(ii) *Indications for use*. For the control of American foulbrood (*Paenibacillus larvae*).

(iii) *Limitations*. The drug should be fed early in the spring or fall and consumed by the bees before the main honey flow begins, to avoid contamination of production honey. Complete treatments at least 4 weeks before main honey flow.

[40 FR 13838, Mar. 27, 1975, as amended at 50 FR 49841, Dec. 5, 1985; 59 FR 14365, Mar. 28, 1994; 62 FR 39443, July 23, 1997; 68 FR 24879, May 9, 2003; 70 FR 69439, Nov. 16, 2005; 73 FR 76946, Dec. 18, 2008; 75 FR 76259, Dec. 8, 2010; 76 FR 59024, Sept. 23, 2011; 77 FR 29217, May 17, 2012]

§ 520.2645 Tylvalosin.

(a) *Specifications*. Granules containing 62.5 percent tylvalosin (w/w) as tylvalosin tartrate.

(b) *Sponsor*. See No. 066916 in §510.600(c) of this chapter.

(c) *Related tolerances*. See §556.748 of this chapter.

(d) *Conditions of use in swine*—(1) *Amount*. Administer 50 parts per million tylvalosin in drinking water for 5 consecutive days.

(2) *Indications for use*. For the control of porcine proliferative enteropathy (PPE) associated with *Lawsonia intracellularis* infection in groups of swine in buildings experiencing an outbreak of PPE.

(3) *Limitations*. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[77 FR 55415, Sept. 10, 2012]

PART 522—IMPLANTATION OR INJECTABLE DOSAGE FORM NEW ANIMAL DRUGS

Sec.

- 522.23 Acepromazine.
- 522.44 Sterile sodium acetazolamide.
- 522.52 Alfaxalone.
- 522.56 Amikacin.
- 522.62 Aminopentamide hydrogen sulfate injection.
- 522.82 Aminopropazine fumarate sterile solution injection.
- 522.84 Beta-aminopropionitrile fumarate.
- 522.88 Sterile amoxicillin trihydrate for suspension.
- 522.90 Ampicillin implantation and injectable dosage forms.
- 522.90a Ampicillin trihydrate sterile suspension.
- 522.90b Ampicillin trihydrate.
- 522.90c Ampicillin sodium.
- 522.144 Arsenamide sodium aqueous injection.
- 522.147 Atipamezole.
- 522.150 Azaperone.
- 522.161 Betamethasone acetate and betamethasone disodium phosphate aqueous suspension.
- 522.163 Betamethasone dipropionate and betamethasone sodium phosphate aqueous suspension.
- 522.204 Boldenone.
- 522.234 Butamisol hydrochloride.
- 522.246 Butorphanol.
- 522.275 N-Butylscopolammonium bromide.
- 522.300 Carfentanil citrate injection.
- 522.304 Carprofen.
- 522.311 Cefovecin.
- 522.313 Ceftiofur injectable dosage forms.
- 522.313a Ceftiofur crystalline free acid.
- 522.313b Ceftiofur hydrochloride.
- 522.313c Ceftiofur sodium.
- 522.380 Chloral hydrate, pentobarbital, and magnesium sulfate sterile aqueous solution.
- 522.390 Chloramphenicol injection.
- 522.460 Cloprostenol sodium.