

§522.1150

21 CFR Ch. I (4-1-13 Edition)

drug to use by or on the order of a licensed veterinarian.

[49 FR 45124, Nov. 15, 1984, as amended at 51 FR 11438, Apr. 3, 1986; 51 FR 25032, July 10, 1986; 53 FR 19773, May 31, 1988; 53 FR 22297, June 15, 1988; 56 FR 50814, Oct. 9, 1991; 57 FR 2837, Jan. 24, 1992; 59 FR 33198, June 28, 1994; 61 FR 59003, Nov. 20, 1996; 63 FR 59216, Nov. 3, 1998; 71 FR 1689, Jan. 11, 2006; 71 FR 39204, July 12, 2006; 75 FR 1274, Jan. 11, 2010; 75 FR 10167, Mar. 5, 2010]

§522.1150 Hydrochlorothiazide injection.

(a) Specifications. Each milliliter contains 25 milligrams of hydrochlorothiazide.

(b) Sponsor. See No. 050604 in §510.600(c) of this chapter.

(c) Conditions of use—(1) Amount. 5 to 10 milliliters (125 to 250 milligrams) intravenously or intramuscularly once or twice a day. After onset of diuresis, treatment may be continued with an orally administered maintenance dose.

(2) Indications for use. For use in cattle as an aid in the treatment of postparturient udder edema.¹

(3) Limitations. Animals should be regularly and carefully observed for early signs of fluid and electrolyte imbalance. Take appropriate countermeasures if this should occur. Milk taken from dairy animals during treatment and for 72 hours (6 milkings) after the latest treatment must not be used for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.¹

[43 FR 59058, Dec. 19, 1978, as amended at 62 FR 63271, Nov. 28, 1997]

§522.1155 Imidocarb dipropionate sterile powder.

(a) Specifications. Imidocarb dipropionate powder is reconstituted with sterile water. Each milliliter of solution contains 100 milligrams of imidocarb base.

(b) Sponsor. No. 000061 in §510.600(c) of this chapter.

(c) Conditions of use. The drug is used in horses and zebras as follows:

¹These conditions are NAS/NRC reviewed and deemed effective. Applications for these uses need not include effectiveness data as specified by §514.111 of this chapter, but may require bioequivalency and safety information.

(1) Amount. For *Babesia caballi* infections, use intramuscularly 2 milligrams of imidocarb base per kilogram of body weight, repeating dosage once after 24 hours. For *Babesia equi* infections, use 4 milligrams of imidocarb base per kilogram of body weight, repeating dosage four times at 72-hour intervals.

(2) Indications for use. For the treatment of babesiosis (piroplasmosis) caused by *Babesia caballi* and *Babesia equi*.

(3) Limitations. Administer intramuscularly in the neck region. Do not inject intravenously. Do not use for other equidae or for animals of other species. Do not use in horses less than 1 year old. Do not use for animals in near-term pregnancies. Imidocarb dipropionate is a cholinesterase inhibitor. Do not use this product simultaneously with or a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides, or chemicals. Do not use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian. Imidocarb dipropionate is sold only under permit issued by the Director of the National Program Planning Staff, Veterinary Services, APHIS, USDA, to licensed or full-time State, Federal, or military veterinarians.

[43 FR 40455, Sept. 12, 1978, as amended at 46 FR 48642, Oct. 2, 1981; 61 FR 8873, Mar. 6, 1996; 62 FR 61625, Nov. 19, 1997]

§522.1156 Imidocarb dipropionate solution.

(a) Specifications. Each milliliter of injectable solution contains 120 milligrams of imidocarb.

(b) Sponsor. See No. 000061 in §510.600(c) of this chapter.

(c) [Reserved]

(d) Conditions of use—(1) Dogs—(i) Amount. 6.6 milligrams imidocarb per kilogram (3 milligrams per pound) of body weight.

(ii) Indications for use. Treatment of clinical signs of babesiosis and/or demonstrated *Babesia* organisms in the blood.

(iii) Limitations. Use subcutaneously or intramuscularly. Not for intravenous use. Repeat the dose after 2 weeks for a total of two treatments.