

Environmental Protection Agency

§ 180.507

§ 180.506 Cyclanilide; tolerances for residues.

(a) *General.* Tolerances are established for residues of the plant growth regulator, cyclanilide, [1-(2,4-dichlorophenylaminocarbonyl)-cyclopropane carboxylic acid] determined as 2,4-dichloroaniline (calculated as cyclanilide) in or on the following food commodities and processed feed:

Commodity	Parts Per Million
Cattle, fat	0.10
Cattle, meat	0.02
Cattle, meat byproducts, except kidney	0.2
Cattle, kidney	2.0
Cotton, undelinted seed	0.60
Cotton, gin byproducts	25.0
Goat, fat	0.10
Goat, meat	0.02
Goat, meat byproducts, except kidney	0.20
Goat, kidney	2.0
Horse, fat	0.10
Horse, meat	0.02
Horse, meat byproducts, except kidney	0.20
Horse, kidney	2.0
Hog, fat	0.10
Hog, meat	0.02
Hog, meat byproducts, except kidney	0.20
Hog, kidney	2.0
Milk	0.04
Sheep, fat	0.10
Sheep, meat	0.20
Sheep, meat byproducts, except kidney	0.20
Sheep, kidney	2.0

(b) *Section 18 emergency exemptions.* [Reserved]

(c) *Tolerances with regional registrations.* [Reserved]

(d) *Indirect or inadvertent residues.* [Reserved]

[62 FR 28355, May 23, 1997; 62 FR 34182, June 25, 1997]

§ 180.507 Azoxystrobin; tolerances for residues.

(a) *General.* (1) Tolerances are established for residues of the fungicide, azoxystrobin, including its metabolites and degradates, in or on the commodities in the following table. Compliance with the tolerance levels specified in the table is to be determined by measuring only the sum of azoxystrobin, [methyl(*E*)-2-(2-(6-(2-cyanophenoxy)pyrimidin-4-yloxy)phenyl)-3-methoxyacrylate], and the *Z*-isomer of azoxystrobin [methyl(*Z*)-2-(2-(6-(2-cyanophenoxy)pyrimidin-4-yloxy)phenyl)-3 methoxyacrylate] in or on the commodity.

Commodity	Parts per million
Acerola	2.0
Almond, hulls	4.0
Animal feed, nongrass, group 18, forage	45
Animal feed, nongrass, group 18, hay	120
Artichoke, globe	4.0
Asparagus	0.04
Atemoya	2.0
Avocado	2.0
Banana	*
Barley, bran	6.0
Barley, forage	25
Barley, grain	3.0
Barley, hay	10.0
Barley, straw	15.0
Berry, low growing, subgroup 13-07G, except cranberry	10.0
Biriba	2.0
Brassica, head and stem, subgroup 5A	3.0
Brassica, leafy greens, subgroup 5B	25
Bushberry subgroup 13-07B	5.0
Caneberry subgroup 13-07A	5.0
Canistel	2.0
Cherimoya	2.0
Cilantro, leaves	30.0
Citrus, dried pulp	20.0
Citrus, oil	40.0
Corn, field, forage	12.0
Corn, field, grain	0.05
Corn, field, refined oil	0.3
Corn, field, stover	25.0
Corn, pop, grain	0.05
Corn, pop, stover	25.0
Corn, sweet, forage	12.0
Corn, sweet, kernel plus cob with husks removed	0.05
Corn, sweet, stover	25.0
Cotton, gin byproducts	45
Cottonseed subgroup 20C	0.7
Cranberry	0.50
Custard apple	2.0
Dragon fruit	2.0
Feijoa	2.0
Fruit, citrus, group 10-10	15.0
Fruit, small vine climbing, except fuzzy kiwifruit, subgroup 13-07F	2.0
Fruit, stone, group 12	1.5
Ginseng ¹	0.5
Grain, aspirated fractions	420
Grass, forage	15
Grass, hay	20
Guava	2.0
Herb Subgroup 19A, dried leaves	260
Herb Subgroup 19A, fresh leaves	50
Hop, dried cones	20.0
Ilama	2.0
Jaboticaba	2.0
Jackfruit	2.0
Longan	2.0
Loquat	2.0
Lychee	2.0
Mango	2.0
Nut, tree, group 14	0.02
Oats, forage	5.0
Oats, grain	1.5
Oats, hay	10.0
Oats, straw	3.0
Onion, bulb, subgroup 3-07A	1.0
Onion, green, subgroup 3-07B	7.5
Papaya	2.0
Passionfruit	2.0
Pawpaw	2.0
Pea and bean, dried shelled, except soybean, subgroup 6C	0.5

§ 180.509

Commodity	Parts per million
Pea and bean, succulent shelled, subgroup 6B	0.5
Peanut	0.2
Peanut, hay	15.0
Peanut, refined oil	0.6
Pepper/eggplant subgroup 8–10B	3.0
Peppermint, tops	30
Persimmon	2.0
Pistachio	0.50
Pulasan	2.0
Rambutan	2.0
Rapeseed subgroup 20A	1.0
Rice, grain	5.0
Rice, hulls	20
Rice, straw	12
Rice, wild, grain	5.0
Rye, forage	7.0
Rye, grain	0.2
Rye, straw	1.5
Sapodilla	2.0
Sapote, black	2.0
Sapote, mamey	2.0
Sapote, white	2.0
Sorghum, grain, forage	25
Sorghum, grain, grain	11
Sorghum, grain, stover	40
Soursop	2.0
Soybean, hay	55.0
Soybean, hulls	1.0
Soybean, seed	0.5
Spanish lime	2.0
Spearmint, tops	30
Spice Subgroup 19B, except black pepper	38
Star apple	2.0
Starfruit	2.0
Sugar apple	2.0
Sugarcane, cane	0.2
Sunflower subgroup 20B	0.5
Tamarind	2.0
Tomato, paste	0.6
Tomato subgroup 8–10A	0.2
Turnip, greens	25
Vegetable, cucurbit, group 9	0.3
Vegetable, foliage of legume, group 7	30.0
Vegetable, leafy, except brassica, group 4	30.0
Vegetable, leaves of root and tuber, group 2	50.0
Vegetable, legume, edible podded, subgroup 6A, except soybean	3.0
Vegetable, root, subgroup 1A	0.5
Vegetable, tuberous and corm, subgroup 1C	8.0
Wasabi, dry	260
Wasabi, fresh	50
Watercress	3.0
Wax jambu	2.0
Wheat, forage	15.0
Wheat, grain	0.2
Wheat, hay	30.0
Wheat, straw	10.0

*2.0 (of which not more than 0.1 is contained in the pulp)
 †There are no United States registrations for use of azoxystrobin on ginseng.

(2) Tolerances are established for residues of the fungicide, azoxystrobin, including its metabolites and degradates, in or on the commodities in the following table. Compliance with the tolerance levels specified in the table is to be determined by measuring only azoxystrobin, [methyl(E)-2-(2-(6-(2-cyanophenoxy) pyrimidin-4-

40 CFR Ch. I (7–1–13 Edition)

yl oxy)phenyl)-3-methoxyacrylate] in or on the commodity.

Commodity	Parts per million
Cattle, fat	0.03
Cattle, meat	0.01
Cattle, meat byproducts	0.07
Goat, fat	0.03
Goat, meat	0.01
Goat, meat byproducts	0.07
Hog, fat	0.010
Hog, meat	0.01
Hog, meat byproducts	0.010
Horse, fat	0.03
Horse, meat	0.01
Horse, meat byproducts	0.07
Milk	0.006
Sheep, fat	0.03
Sheep, meat	0.01
Sheep, meat byproducts	0.07

(b) *Section 18 emergency exemptions.*
 [Reserved]

(c) *Tolerances with regional registration.* [Reserved]

(d) *Indirect or inadvertent residues.*
 [Reserved]

[62 FR 32235, June 13, 1997]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 180.507, see the List of CFR Sections Affected, which appears in the Finding Aids section of the printed volume and at www.fdsys.gov.

§ 180.509 Mefenpyr-diethyl; tolerance for residues.

(a) *General.* Tolerances are established for residues of the safener, mefenpyr-diethyl, including its metabolites and degradates, when applied at a rate no greater than 0.053 pound safener per acre per growing season in or on the commodities in the table below. Compliance with the tolerance levels specified below is to be determined by measuring only the sum of mefenpyr-diethyl (1-(2,4-dichlorophenyl)-4,5-dihydro-5-methyl-1H-pyrazole-3,5-dicarboxylic acid, diethyl ester) and its 2,4-dichlorophenyl-pyrazoline metabolites, calculated as the stoichiometric equivalent of mefenpyr-diethyl, in or on the commodity.

Commodity	Parts per million
Barley, grain	0.05
Barley, hay	0.2
Barley, straw	0.5
Cattle, meat byproducts	0.1
Goat, meat byproducts	0.1
Grass, forage	1.6