Environmental Protection Agency

- (b) Section 18 emergency exemptions. [Reserved]
- (c) Tolerances with regional registrations. A tolerance with regional registration as defined in §180.1(1) is established for residues of the fungicide fenamidone, including its metabolites and degradates, in or on the following commodities. Compliance with the tolerance levels is to be determined by measuring only fenamidone (4H-Imidazol-4-one, 3,5-dihydro-5-methyl-2-(methylthio)-5-phenyl-3 (phenylamino)-,(S)-), in or on the commodity:

| Commodity | Parts per million |
|--------------------|----------------------|
| Grape ¹ | 1.0 |

¹ Applicable to grapes grown East of the Rocky Mountains.

(d) Indirect or inadvertent residues. Tolerances are established for residues of the fungicide fenamidone, including its metabolites and degradates, in or on the following commodities. Compliance with the tolerance levels is to be determined by measuring fenamidone (4H-Imidazol-4-one, 3,5-dihydro-5-methyl-2-(methylthio)-5-phenyl-3

(phenylamino)-,(S)-), and its metabolite RPA 717879 (2,4-imidazolidinedione, 5-methyl-5-phenyl), in or on the following commodities when present therein as a result of application of fenamidone to the crops in paragraph (a)(1).

| Commodity | Parts per million |
|--------------------------------------|----------------------|
| Grain, cereal, group 15, except rice | 0.1 |
| 16, except rice | 0.5 |
| Soybean, forage | 0.15 |
| Soybean, hay | 0.25 |
| Soybean, seed | 0.02 |
| Strawberry | 0.02 |

[67 FR 60976, Sept. 27, 2002, as amended at 69 FR 58066, Sept. 29, 2004; 71 FR 55293, Sept. 22, 2006; 72 FR 60272, Oct. 24, 2007; 74 FR 34257, July 15, 2009; 76 FR 34885, June 15, 2011; 76 FR 70895, Nov. 16, 2011; 77 FR 32401, June 1, 2012]

§ 180.580 Iodosulfuron-Methyl-Sodium; tolerances for residues.

(a) General. Tolerances are established for residues of the herbicide Iodosulfuron-Methyl-Sodium (methyl 4-iodo-2-[3-(4-methoxy-6-methyl-1,3,5 trizzin-2-yl)ureidosulfonyl]benzoate, sodium salt) in or on the following commodities:

| Commodity | Parts per million |
|---------------------|----------------------|
| Corn, field, forage | 0.05 |
| Corn, field, grain | 0.03 |
| Corn, field, stover | 0.05 |
| Wheat, forage | 0.10 |
| Wheat, grain | 0.02 |
| Wheat, hay | 0.05 |
| Wheat, straw | 0.05 |

- (b) Section 18 emergency exemptions. [Reserved]
- (c) Tolerances with regional registrations. [Reserved]
- (d) Indirect or inadvertent residues. [Reserved]

[67 FR 57532, Sept. 11, 2002, as amended at 74 FR 23644, May 20, 2009]

§ 180.581 Iprovalicarb; tolerances for residues.

(a) General. Tolerances are established for residues of iprovalicarb, [2-methyl-1[[[(1S)-(4-methylphenyl) ethyl] amino]carbonyl] propyl]carbamic acid methylethylester, in or on the following commodities.

| Commodity | Parts per million |
|--------------------|----------------------|
| Grape ¹ | 2.0 1.0 |

¹There is no U.S. registration as of September 1, 2005.

- (b) Section 18 emergency exemptions. [Reserved]
- (c) Tolerances with regional registrations. [Reserved]
- (d) Indirect or inadvertent residues. [Reserved]

[67 FR 54359, Aug. 22, 2002, as amended at 70 FR 55281, Sept. 21, 2005]

§ 180.582 Pyraclostrobin; tolerances for residues.

(a) General. (1) Tolerances are established for residues of the fungicide pyradostrobin, including its metabolites and degradates, in or on the commodities in the table below. Compliance with the tolerance levels specified below is to be determined by measuring only the sum of pyraclostrobin (carbamic acid, [2-[[[1-(4-chlorophenyl)-1H-pyrazol-3-yl]oxy] methyl]phenyl]methoxy-, methyl ester) and its desmethoxy metabolite (methyl-N-[[[1-(4-chlorophenyl)-1H-pyrazol-3-