

## §51.1016

(d) Scars which are dark, rough, or deep and aggregate more than 5 percent of the fruit surface, or scars which are fairly light in color, slightly rough, or of slight depth and aggregate more than 10 percent of the fruit surface, or scars which are light colored, fairly smooth, with no depth and aggregate more than 25 percent of the fruit surface;

(e) Thorn scratches when the injury is not well healed, or when dark colored, rough or deep and aggregates more than 5 percent of the fruit surface, or when light colored, fairly smooth and concentrated and aggregates more than 10 percent of the fruit surface, or light colored and scattered thorn injury which detracts from the appearance of the fruit to a greater extent than the 10 percent light colored concentrated injury;

(f) Scale when the appearance of the fruit is affected to a greater extent than that of a lime which has a blotch the area of a circle one-half inch in diameter;

(g) Sunburn which causes decided flattening of the fruit, marked drying or dark discoloration of the skin, material drying of the flesh underneath the affected area, or which affects more than 10 percent of the fruit surface;

(h) Scab which seriously affects shape or texture;

(i) Blanching when more than 50 percent, in the aggregate, of the fruit surface shows a whitish to yellowish green area or areas because of shading, resting on the surface of the ground, or contact with other fruit on the tree. Such areas are not to be confused with limes which are turning yellow due to the ripening process;

(j) Yellow color when plainly visible and caused by the ripening process;

(k) Discoloration caused by rust mite, melanose or other means, when fairly smooth and more than 50 percent of the fruit surface is affected, or when slightly rough and more than 25 percent of the fruit surface is affected; and,

(l) Buckskin when more unsightly than the maximum discoloration allowed, or the fruit texture is seriously affected.

## 7 CFR Ch. I (1-1-13 Edition)

### §51.1016 Diameter.

*Diameter* means the greatest dimension measured at right angles to a line from stem to blossom end of the fruit.

### Subpart—United States Standards for Grades of Florida Oranges and Tangelos

SOURCE: 61 FR 20708, May 8, 1996, unless otherwise noted.

#### GENERAL

### §51.1140 General.

The standards contained in this subpart apply only to the common or sweet orange group and varieties and hybrids of varieties belonging to the Mandarin group, except tangerines, and to the citrus fruit commonly known as “tangelo”—a hybrid between tangerine or mandarin orange (*Citrus reticulata*) with either the grapefruit or pomelo (*C. paradisi* and *C. grandis*). Separate U.S. standards apply to tangerines. The standards for internal quality contained in §§51.1176 through 51.1179 apply only to common sweet oranges (*Citrus sinensis* (L) Osbeck).

#### GRADES

### §51.1141 U.S. Fancy.

“U.S. Fancy” consists of oranges which meet the following requirements:

(a) Basic requirements:

(1) Discoloration: Not more than one-tenth of the surface, in the aggregate, may be affected by discoloration. (See §51.1161.);

(2) Firm;

(3) Mature;

(4) Similar varietal characteristics;

(5) Smooth texture;

(6) Well colored; and,

(7) Well formed.

(b) Free from:

(1) Ammoniation;

(2) Buckskin;

(3) Caked melanose;

(4) Creasing;

(5) Decay;

(6) Scab;

(7) Split navels;

(8) Sprayburn;

(9) Undeveloped segments;

(10) Unhealed skin breaks; and,