

§ 201.57

(ii) Less than half of the original cotyledon tissue free of necrosis or decay.

(2) Epicotyl:

(i) Missing. (May be assumed to be present if the cotyledons are intact.)

(ii) [Reserved]

(3) Hypocotyl:

(i) Malformed, such as markedly shortened, curled, or thickened.

(ii) Deep open cracks extending into the conducting tissue.

(iii) Watery.

(4) Root:

(i) None.

(ii) Missing or stubby primary root with weak secondary or adventitious roots.

(5) Seedling:

(i) One or more essential structures impaired as a result of decay from primary infection.

(ii) Albino.

[59 FR 64506, Dec. 14, 1994]

§ 201.57 Hard seeds.

Seeds which remain hard at the end of the prescribed test because they have not absorbed water, due to an impermeable seed coat, are to be counted as "hard seed." If at the end of the germination period provided for legumes, okra, cotton and dichondra in these rules and regulations there are still present swollen seeds or seeds of these kinds which have just started to germinate, all seeds or seedlings except the above-stated shall be removed and the test continued for 5 additional days and the normal seedlings included in the percentage of germination. For flatpea, continue the swollen seed in test for 14 days when germinating at 15–25 °C or for 10 days when germinating at 20 °C.

[5 FR 33, Jan. 4, 1940, as amended at 10 FR 9952, Aug. 11, 1945; 20 FR 7936, Oct. 21, 1955; 65 FR 1708, Jan. 11, 2000]

§ 201.57a Dormant seeds.

Dormant seeds are viable seeds, other than hard seeds, which fail to germinate when provided the specified germination conditions for the kind of seed in question.

(a) Viability of ungerminated seeds shall be determined by any of the following methods or combinations of methods: a cutting test, tetrazolium

7 CFR Ch. I (1–1–13 Edition)

test, scarification, or application of germination promoting chemicals.

(b) The percentage of dormant seed, if present, shall be determined in addition to the percentage of germination for the following kinds: Bahiagrass, basin wildrye, big bluestem, little bluestem, sand bluestem, yellow bluestem, bottlebrush-squirreltail, buffalograss, buffelgrass, galletagrass, forage kochia, blue grama, side-oats grama, Indian ricegrass, johnsongrass, sand lovegrass, weeping lovegrass, mountain rye, sand dropseed, smilo, switchgrass, veldtgrass, western wheatgrass, and yellow indiagrass.

(c) For green needlegrass, if the test result of method 2 is less than the result of method 1, subtract the result of method 2 from method 1 and report the difference as the percentage of dormant seed. Refer to § 201.58(b)(7).

[46 FR 53638, Oct. 29, 1981, as amended at 59 FR 64506, Dec. 14, 1994]

§ 201.58 Substrata, temperature, duration of test, and certain other specific directions for testing for germination and hard seed.

Specific germination requirements are set forth in table 2 to which the following paragraphs (a), (b), and (c) are applicable.

(a) *Definitions and explanations applicable to table 2—(1) Duration of tests.* The following deviations are permitted from the specified duration of tests: Any test may be terminated prior to the number of days listed under "Final count" if the maximum germination of the sample has then been determined. The number of days stated for the first count is approximate and a deviation of 1 to 3 days is permitted. If at the time of the prescribed test period the seedlings are not sufficiently developed for positive evaluation, it is possible to extend the time of the test period two additional days. (Also, see paragraph (a)(5) of this section and 201.57.)

(2) *Light.* Cool white fluorescent light shall be provided where light is required in table 2. The light intensity shall be 75 to 125 foot-candles (750–1,250 lux). (The light intensity for nondormant seed and during seedling development may be as low as 25 foot-candles to enable the essential structures to be evaluated with greater certainty.) The