### § 520.606

tranquilizers, or central nervous system depressants.

- (4) Do not use in horses which are severly debilitated or suffering from diarrhea or severe constipation, infectious disease, toxemia, or colic. Do not administer to horses affected with chronic alveolar emphysema (heaves) or other respiratory conditions.
- (5) Do not use in horses intended for food purposes.
- (6) Federal law restricts this drug to use by or on the order of a licensed veterinarian.
- (i) Conditions of use in dogs, cats, puppies, and kittens. (1) Each tablet contains 2, 5, 10, or 20 milligrams of dichloryos.
- (2) It is administered orally at 5 milligrams of dichlorvos per pound of body weight.
- (3) Dogs and puppies: Removal and control of intestinal roundworms (*Toxocara canis* and *Toxascaris leonina*) and hookworms (*Ancylostoma caninum* and *Uncinaria stenocephala*).
- (4) Cats and kittens: Removal and control of intestinal roundworms (Toxocara cati and Toxascaris leonina) and hookworms (Ancylostoma tubaeforme and Uncinaria stenocephala).
- (5) Dichlorvos is a cholinesterase inhibitor. Do not use simultaneously with or within a few days before or after treatment with or exposure to cholinesterase-inhibiting drugs, pesticides, or chemicals.
- (6) Do not use in animals under 10 days of age or 1 pound of body weight.
- (7) Do not administer to animals showing signs of constipation, mechanical blockage of the intestinal tract, impaired liver function, or recently exposed to or showing signs of infectious disease.
- (8) Do not use in dogs or puppies infected with *Dirofilaria immitis*.
- (9) Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[40 FR 13838, Mar. 27, 1975, as amended at 48 FR 40704, Sept. 9, 1983; 51 FR 28546, Aug. 8, 1986; 62 FR 35076, June 30, 1997; 64 FR 18571, Apr. 15, 1999; 78 FR 21059, Apr. 9, 2013]

#### § 520.606 Diclazuril.

(a) Specifications. Each 100 grams (g) of pellets contain 1.56 g diclazuril.

- (b) Sponsor. See No. 000061 in §510.600(c) of this chapter.
- (c) Conditions of use in horses(1) Amount. Administer 1 milligram (mg) per kilogram (0.45 mg per pound) of body weight in the daily grain ration for 28 days.
- (2) Indications for use. For the treatment of equine protozoal myeloencephalitis (EPM) caused by Sarcocystis neurona.
- (3) *Limitations*. Do not use in horses intended for human consumption. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[72 FR 20943, Apr. 27, 2007]

## § 520.608 Dicloxacillin sodium monohydrate capsules.

- (a) Specifications. Each capsule contains dicloxacillin sodium monohydrate equivalent to 50, 100, 200, or 500 milligrams of dicloxacillin.
- (b) Sponsor. See No. 000856 in §510.600 (c) of this chapter.
- (c) Conditions of use. Dogs(1) Amount. 5 to 10 milligrams per pound of body weight, three times daily. In severe cases, up to 25 milligrams per pound of body weight three times daily.
- (2) Indications for use. Treatment of pyoderma (pyogenic dermatitis) due to penicillinase-producing staphylococci sensitive to the drug.
- (3) Limitations. For the treatment of dogs only. Continue treatment for 24 to 48 hours after the animal has become afebrile or asymptomatic, Administer 1 to 2 hours before feeding to ensure maximum absorption. Not for use in animals which are raised for food production. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

[57 FR 37325, Aug. 18, 1992]

# § 520.620 Diethylcarbamazine oral dosage forms.

# § 520.622 Diethylcarbamazine citrate oral dosage forms.

## § 520.622a Diethylcarbamazine citrate tablets.

(a) Sponsors. (1) [Reserved]

(2) See 053501 in §510.600(c) of this chapter for use of 100, 200, and 300 milligram tablets for prevention of heartworm disease in dogs and as an aid in