

(4) Federal law restricts this drug to use by or on the order of a licensed veterinarian.¹

[40 FR 13838, Mar. 27, 1975, as amended at 46 FR 48642, Oct. 2, 1981; 61 FR 8873, Mar. 6, 1996; 62 FR 61624, Nov. 19, 1997]

§ 520.870 Etodolac.

(a) *Specifications.* Each tablet contains 150, 300, or 500 milligrams (mg) of etodolac.

(b) *Sponsor.* See No. 000010 in § 510.600(c) of this chapter.

(c) [Reserved]

(d) *Conditions of use*—(i) *Amount.* Administer 10 to 15 mg per kilogram (4.5 to 6.8 mg per pound) of body weight per day orally.

(ii) *Indications for use.* For the management of pain and inflammation associated with osteoarthritis in dogs.

(iii) *Limitations.* Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(2) [Reserved]

[63 FR 51300, Sept. 25, 1998, as amended at 68 FR 51705, Aug. 28, 2003; 75 FR 10166, Mar. 5, 2010]

§ 520.903 Febantel oral dosage forms.

§ 520.903a Febantel paste.

(a) *Chemical name.* Dimethyl [[2-[(methoxyacetyl)amino]-4-(phenylthio)phenyl] carbonimidoyl]bis [carbamate].

(b) *Specifications.* The drug is a paste containing 45.5 percent febantel.

(c) *Sponsor.* See No. 000859 in § 510.600(c) of this chapter.

(d) *Conditions of use*—(1) *Amount.* Six milligrams per kilogram (2.73 milligrams per pound) of body weight in horses.

(2) *Indications for use.* For removal of large strongyles (*Strongylus vulgaris*, *S. edentatus*, *S. equinus*); ascarids (*Parascaris equorum*—sexually mature and immature); pinworms (*Oxyuris equi*—adult and 4th stage larva); and the various small strongyles in horses, foals, and ponies.

(3) *Limitations.* (i) The paste may be administered on the base of the tongue or well mixed into a portion of the normal grain ration.

(ii) [Reserved]

(iii) For animals maintained on premises where reinfection is likely to

occur, retreatment may be necessary. For most effective results, retreat in 6 to 8 weeks.

(iv) Not for use in horses intended for food.

(v) Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism.

[43 FR 8797, Mar. 3, 1978; 43 FR 12311, Mar. 24, 1978, as amended at 43 FR 60882, Dec. 29, 1978. Redesignated at 45 FR 8587, Feb. 8, 1980]

§ 520.903b Febantel suspension.

(a) *Specifications.* The suspension contains 9.3 percent (2.75 grams per ounce) febantel.

(b) *Sponsor.* See 000859 in § 510.600(c) of this chapter.

(c) *Conditions of use*—(1) *Amount.* 3 milliliters per 100 pounds body weight or 1 fluid ounce per 1000 pounds (6 milligrams per kilogram body weight).

(2) *Indications for use.* For removal of ascarids (*Parascaris equorum*—adult and sexually immature), pinworms (*Oxyuris equi*—adult and 4th stage larvae), large strongyles (*Strongylus vulgaris*, *S. edentatus*, *S. equinus*), and the various small strongyles in horses, breeding stallions and mares, pregnant mares, foals, and ponies.

(3) *Limitations.* Administer by stomach tube or drench, or by mixing well into a portion of the normal grain ration. For animals maintained on premises where reinfection is likely to occur, retreatment may be necessary. For most effective results, retreat in 6 to 8 weeks. Not for use in horses intended for food. Federal law restricts this drug to use by or on the order of a licensed veterinarian.

(d) *Special considerations.* Febantel suspension may be used in combination with trichlorfon oral liquid in accordance with the provisions of § 520.2520c, this section, and the following conditions:

(1) Combine 1 part febantel suspension with 5 parts trichlorfon liquid.

(2) Allow animal to consume a portion of daily grain ration; administer mixture by stomach tube at rate of 18 milliliters per 100 pounds of body weight.

[45 FR 8587, Feb. 8, 1980]