

Food and Drug Administration, HHS

§ 556.460

12.5 micrograms per kilogram of body weight per day.

(b) *Tolerances.* The tolerances for residues of monensin are:

(1) *Cattle*—(i) *Liver.* 0.10 part per million (ppm).

(ii) *Muscle, kidney, and fat.* 0.05 ppm.

(iii) *Milk.* Not required.

(2) *Goats*—(i) *Edible tissues.* 0.05 ppm.

(ii) [Reserved]

(3) *Chickens, turkeys, and quail.* A tolerance for residues of monensin in chickens, turkeys, and quail is not required.

(c) *Related conditions of use.* See §§ 520.1448 and 558.355 of this chapter.

[64 FR 5159, Feb. 3, 1999, as amended at 69 FR 68783, Nov. 26, 2004; 72 FR 56897, Oct. 5, 2007]

§ 556.425 Morantel tartrate.

A tolerance of 0.7 part per million is established for N-methyl-1,3-propanediamine (MAPA, marker residue) in the liver (target tissue) of cattle and goats. A tolerance for residues of morantel tartrate in milk is not required.

[59 FR 17922, Apr. 15, 1994]

§ 556.426 Moxidectin.

(a) *Acceptable daily intake (ADI).* The ADI for total residues of moxidectin is 4 micrograms per kilogram of body weight per day.

(b) *Tolerances*—(1) *Cattle*—(i) *Fat (the target tissue).* The tolerance for parent moxidectin (the marker residue) is 900 parts per billion (ppb).

(ii) *Liver.* The tolerance for parent moxidectin (the marker residue) is 200 ppb.

(iii) *Muscle.* The tolerance for parent moxidectin (the marker residue) is 50 ppb.

(iv) *Milk.* The tolerance for parent moxidectin (the marker residue) is 40 ppb.

(2) *Sheep*—(i) *Fat (the target tissue).* The tolerance for parent moxidectin (the marker residue) is 900 parts per billion (ppb).

(ii) *Liver.* The tolerance for parent moxidectin (the marker residue) is 200 ppb.

(iii) *Muscle.* The tolerance for parent moxidectin (the marker residue) is 50 ppb.

(c) *Related conditions of use.* See §§ 520.1454 and 522.1450 of this chapter.

[65 FR 36617, June 9, 2000, as amended at 65 FR 76930, Dec. 8, 2000; 70 FR 36338, June 23, 2005; 70 FR 76163, Dec. 23, 2005]

§ 556.428 Narasin.

(a) *Acceptable daily intake (ADI).* The ADI for total residues of narasin is 5 micrograms per kilogram of body weight per day.

(b) *Tolerances*—(1) *Chickens (abdominal fat).* The tolerance for parent narasin (the marker residue) is 480 parts per billion.

(2) [Reserved]

[66 FR 23589, May 9, 2001]

§ 556.430 Neomycin.

(a) *Acceptable daily intake (ADI).* The ADI for total residues of neomycin is 6 micrograms per kilogram of body weight per day.

(b) *Tolerances.* Tolerances are established for residues of parent neomycin in uncooked edible tissues as follows:

(1) *Cattle, swine, sheep, and goats.* 7.2 parts per million (ppm) in kidney (target tissue) and fat, 3.6 ppm in liver, and 1.2 ppm in muscle.

(2) *Turkeys.* 7.2 ppm in skin with adhering fat, 3.6 ppm in liver, and 1.2 ppm in muscle.

(3) *Milk.* A tolerance is established for residues of parent neomycin of 0.15 ppm.

[64 FR 31498, June 11, 1999]

§ 556.440 Nequinatate.

A tolerance of 0.1 part per million is established for negligible residues of nequinatate in the uncooked edible tissues of chickens.

§ 556.445 Nicarbazine.

A tolerance of 4 parts per million is established for residues of nicarbazine in uncooked chicken muscle, liver, skin, and kidney.

[42 FR 56729, Oct. 28, 1977]

§ 556.460 Novobiocin.

Tolerances for residues of novobiocin are established at 0.1 part per million in milk from dairy animals and 1 part