

## § 640.92

value 5 percent for the protein concentration of the product.

(g) *Incubation.* All final containers of Plasma Protein Fraction (Human) shall be incubated at 20 to 35 °C for at least 14 days following the heat treatment prescribed in paragraph (e) of this section. At the end of this incubation period, each final container shall be examined and all containers showing any indication of turbidity or microbial contamination shall not be issued. The contents of turbid final containers shall be examined microscopically and tested for sterility. If growth occurs, the types of organisms shall be identified as to genus and the material from such containers shall not be used for further manufacturing.

[42 FR 27583, May 31, 1977, as amended at 64 FR 26286, May 14, 1999]

### § 640.92 Tests on final product.

Tests shall be performed on the final product to determine that it meets the following standards:

(a) *Protein concentration.* The final product shall be a 5.0 ±0.30 percent solution of protein.

(b) *Protein composition.* The total protein in the final product shall consist of at least 83 percent albumin, and no more than 17 percent globulins. No more than 1 percent of the total protein shall be gamma globulin. The protein composition shall be determined by a method that has been approved for each manufacturer by the Director, Center for Biologics Evaluation and Research, Food and Drug Administration.

(c) *pH.* The pH shall be 7.0 ±0.3 when measured in a solution of the final product diluted to a concentration of 1 percent protein with 0.15 molar sodium chloride.

(d) *Sodium concentration.* The sodium concentration of the final product shall be 130 to 160 milliequivalents per liter.

(e) *Potassium concentration.* The potassium concentration of the final product shall not exceed 2 milliequivalents per liter.

(f) *Heat stability.* A final container sample of Plasma Protein Fraction (Human) shall remain unchanged, as determined by visual inspection, after heating at 57 °C for 50 hours, when compared to its control consisting of a

## 21 CFR Ch. I (4–1–14 Edition)

sample, from the same lot, which has not undergone this heating.

[42 FR 27583, May 31, 1977, as amended at 49 FR 23834, June 8, 1984; 55 FR 11013, Mar. 26, 1990; 64 FR 26286, May 14, 1999; 65 FR 13679, Mar. 14, 2000]

### § 640.93 General requirements.

(a) *Preservative.* The final product shall not contain a preservative.

(b) *Storage of bulk solution.* After all processing steps have been completed, the sterile bulk solution shall be stored in a manner that will ensure the continued sterility of the product, and at a temperature that shall not exceed the recommended storage temperature of the final product prescribed in §610.53 of this chapter.

### § 640.94 Labeling.

In addition to the labeling requirements of §§610.60, 610.61, and 610.62 of this chapter, the container and package labels shall contain the following information:

(a) The osmotic equivalent in terms of plasma, and the sodium concentration in terms of a value or a range in milliequivalents per liter.

(b) The cautionary statement placed in a prominent position on the label, "Do Not Use if Turbid. Do Not Begin Administration More than 4 Hours After the Container Has Been Entered."

[42 FR 27583, May 31, 1977, as amended at 49 FR 2244, Jan. 19, 1984; 64 FR 26286, May 14, 1999]

## Subpart J—Immune Globulin (Human)

### § 640.100 Immune Globulin (Human).

(a) *Proper name and definition.* The proper name of this product shall be Immune Globulin (Human). The product is defined as a sterile solution containing antibodies derived from human plasma.

(b) *Source material.* The source material of Immune Globulin (Human) shall be plasma recovered from Whole Blood prepared as prescribed in §§640.1 through 640.5, or Source Plasma prepared as prescribed in §§640.60 through 640.76.