Environmental Protection Agency

Commodity	Parts per million
Almond, hulls	7.0
Apple, wet pomace	4.5
Aspirated grain fractions	95
Banana ¹	0.2
Barley, grain	0.1
Barley, hay	0.05
Barley, straw	0.05
Beet, sugar	0.3
Beet, sugar, dried pulp	1.9
Berry, low growing, subgroup 13-07G, except	
cranberry	2.5
Brassica, head and stem, subgroup 5A	1.9
Brassica, leafy green, subgroup 5B	35
Canola, seed	0.01
Carrot	0.50
Chickpea	0.08
Citrus, dried pulp	2.0
Citrus, oil	25
Corn, sweet, forage	0.01
Corn, sweet, kernel plus cob with husks re-	0.01
moved Corn, sweet, stover	0.01
	0.05
Cotton, gin byproducts	0.05
Cotton, undelinted seed	0.60
Fruit, citrus, group 10–10Fruit, pome, group 11–10	1.0
	2.5
Fruits, stone, group 12	4.0
Grape	6.0
Grape, raisin	0.07
Mango ¹ Nut, tree, group 14	0.07
	0.03
Oat, forage	
Oat, grain	0.01
Oat, hay	
Oat, straw	0.05
Onion, bulb, subgroup 3-07A	0.20
Onion, green, subgroup 3-07B	6.0
Papaya ¹	0.30
Pistachio	0.03
Potato, wet peel	7.3
Rye, forage	0.15
Rye, grain	0.01
Rye, straw	0.05
Soybean, hulls	0.20
Soybean, seed	0.15
Turnip, greens	35
Vegetable, cucurbit, group 9	0.70
Vegetable, fruiting, group 8–10	0.60
Vegetable, tuberous and corm, subgroup 1C	4.0
Wax jambu 1	1.5
Wheat, forage	0.1
Wheat, grain	0.1
Wheat, hay	0.05
Wheat, straw	0.1

¹There are no U.S. registrations.

(2) Tolerances are established for residues of difenoconazole, including its metabolites and degradates, in the commodities in the table below. Compliance with the tolerance levels specified below is to be determined by measuring the sum of difenoconazole, 1-[2-[2-chloro-4-(4-chlorophenoxy)phenyl]-4-methyl-1,3-dioxolan-2-ylmethyl]-1H-1,2,4-triazole, and its metabolite, CGA-205375, 1-[2-chloro-4-(4-chlorophenoxy)phenyl]-2-[1,2,4]triazol-1-ylethanol, calculated as the stoichio-

metric equivalent of difenoconazole, in the following commodities:

Commodity	Parts per million
Cattle, fat	0.10
Cattle, liver	0.40
Cattle, meat	0.05
Cattle, meat byproduct (except liver)	0.10
Egg	0.02
Goat, fat	0.10
Goat, liver	0.40
Goat, meat	0.05
Goat, meat byproduct (except liver)	0.10
Hog, fat	0.10
Hog, liver	0.40
Hog, meat	0.05
Hog, meat byproduct (except liver)	0.10
Horse, fat	0.10
Horse, liver	0.40
Horse, meat	0.05
Horse, meat byproduct (except liver)	0.10
Milk	0.01
Sheep, fat	0.10
Sheep, liver	0.40
Sheep, meat	0.05
Sheep, meat byproduct (except liver)	0.10

- (b) Section 18 emergency exemptions. [Reserved]
- (c) Tolerances with regional registrations. [Reserved]
- (d) Indirect or inadvertent residues. [Reserved]

[64 FR 36254, July 6, 1999]

EDITORIAL NOTE: For FEDERAL REGISTER citations affecting § 180.475, see the List of CFR. Sections Affected, which appears in the Finding Aids section of the printed volume and at www.fdsys.gov.

§ 180.476 Triflumizole; tolerances for residues.

(a) General. (1) Tolerances are established for residues of the fungicide triflumizole, including its metabolites and degradates, in or on the commodities listed in the table below. Compliance with the tolerance levels specified below is to be determined by measuring only the parent compound triflumizole, 1-(1-((4-chloro-2-

(trifluoromethyl)phenyl)imino)-2-propoxyethyl)-1 H -imidazole, and its metabolites containing the 4-chloro-2-trifluoromethylaniline moiety, calculated as stoichiometric equivalent of the parent compound.

Commodity	Parts per million
Apple, dry pomace	2.0 2.0
cranberry	2.0

§ 180.477

Commodity	Parts per million
Brassica, head and stem, subgroup 5A	8.0
Brassica, leafy greens, subgroup 5B	40
Canistel	2.5
Cherry, sweet	1.5
Cherry, tart	1.5
Cilantro, leaves	35
Fruit, pome, group 11-10	0.50
Fruit, small, vine climbing, except fuzzy	
kiwifruit, subgroup 13-07F	2.5
Grape, dried pomace	15.0
Grape, raisin, waste	10.0
Grape, wet pomace	15.0
Hazelnut	0.05
Hop, dried cones	50
Leafy greens subgroup 4A, except spinach	35
Mango	2.5
Papaya	2.5
Pineapple	4.0
Sapodilla	2.5
Sapote, black	2.5
Sapote, mamey	2.5
Star apple	2.5
Swiss chard	18
Tomato	1.5
Turnip, greens	40
Vegetable, cucurbit, Group 9	0.5

(2) Tolerances are established for residues of the fungicide triflumizole, including its metabolites and degradates, in or on the commodities of animal origin listed in the table below. Compliance with the tolerance levels specified below is to be determined by measuring only the parent compound triflumizole, 1-(1-((4-chloro-2-

(trifluoromethyl)phenyl)imino)-2-propoxyethyl)-1 H -imidazole, the metabolite 4-chloro-2-hydroxy-6-trifluoromethylaniline sulfate, and other metabolites containing the 4-chloro-2-trifluoromethylaniline moiety, calculated as the parent compound.

Commodity	Parts per million
Cattle, fat	0.5
Cattle, meat	0.05
Cattle, meat byproducts	0.5
Egg	0.05
Goat, fat	0.5
Goat, meat	0.05
Goat, meat byproducts	0.5
Hog, fat	0.5
Hog, meat	0.05
Hog, meat byproducts	0.5
Horse, fat	0.5
Horse, meat	0.05
Horse, meat byproducts	0.5
Milk	0.05
Poultry, fat	0.05
Poultry, meat	0.05
Poultry, meat byproducts	0.1
Sheep, fat	0.5
Sheep, meat	0.05
Sheep, meat byproducts	0.5

- (b) Section 18 emergency exemptions. [Reserved]
- (c) Tolerances with regional registrations. [Reserved]
- (d) Indirect or inadvertent residues. [Reserved]

[65 FR 33702, May 24, 2000, as amended at 67 FR 40228, June 12, 2002; 67 FR 54587, Aug. 23, 2002; 70 FR 7047, Feb. 10, 2005; 70 FR 17915, Apr. 8, 2005; 71 FR 13279, Mar. 15, 2006; 71 FR 49358, Aug. 23, 2006; 74 FR 26543, June 3, 2009; 74 FR 46376, Sept. 9, 2009; 76 FR 34885, June 15, 2011; 79 FR 12408, Mar. 5, 2014]

§ 180.477 Flumiclorac pentyl; tolerances for residues.

(a) General. Tolerances are established for residues of the herbicide flumiclorac pentyl, including its metabolites and degradates, in or on the commodities in the table in this paragraph. Compliance with the tolerance levels specified in this paragraph is to be determined by measuring only flumiclorac pentyl, pentyl(2-chloro-4-fluoro-5-(1,3,4,5,6,7-hexahydro-1,3-dioxo-2*H*-isoindol-2-yl)phenoxy)acetate, in or on the commodity.

Parts per million
0.01
0.01
0.01
3.0
0.2
0.02
0.01

- (b) Section 18 emergency exemptions. [Reserved]
- (c) Tolerances with regional registrations. [Reserved]
- (d) Indirect or inadvertent residues. [Reserved]

[65 FR 33702, May 24, 2000, as amended at 71 FR 11533, Mar. 8, 2006; 76 FR 23497, Apr. 27, 2011]

§180.478 Rimsulfuron; tolerances for residues.

(a) General. Tolerances are established for residues of the herbicide rimsulfuron, including its metabolites and degradates, in or on the commodities in the table below. Compliance with the tolerance levels specified below is to be determined by measuring only rimsulfuron, N-((4,6-dimethoxypyrimidin-2-yl)aminocarbonyl)-3-(ethylsulfonyl)-2-