

Fishery Conservation and Management

Pt. 660, Subpt. C, Table 1b

TABLE 1b TO PART 660, SUBPART C—2013, ALLOCATIONS BY SPECIES OR SPECIES GROUP (WEIGHTS IN METRIC TONS)

Table 1b. To Part 660, Subpart C - 2013, Allocations by Species or Species Group. (Weights in Metric Tons)

Species	Fishery HG	Allocations			
		Trawl		Non-trawl	
		%	Mt	%	Mt
Arrowtooth flounder	4,070	95%	3,866	5%	203
Bocaccio - S of 40°10' N. lat. a/	311.6	NA	74.9	NA	236.7
Canary rockfish a/ b/	98.5	NA	52.5	NA	46.0
Chilipepper - S of 40°10 N. Lat.	1,466	75%	1,100	25%	367
Cowcod - S of 40°10' N. lat. a/	2.9	NA	1.0	NA	1.9
Darkblotched rockfish c/	296.2	95%	281.4	5%	14.8
Dover sole	23,410	95%	22,240	5%	1,171
English sole	6,712	95%	6,376	5%	336
Lingcod					
N of 40°10' N. lat.	2,758	45%	1,241	55%	1,517
S of 40°10' N. lat.	1,102	45%	496	55%	606
Longnose Skate a/	1,928	90%	1,735	10%	193
Longspine thornyhead					
N of 34°27' N. lat.	1,963	95%	1,865	5%	98
Minor shelf rockfish north a/	903	60.2%	543	39.8%	359
Minor shelf rockfish south a/	668	12.2%	81	87.8%	587
Minor slope rockfish north	1,098	81%	889	19%	209
Minor slope rockfish south	597	63%	376	37%	221
Other flatfish	4,682	90%	4,214	10%	468
Pacific cod	1,191	95%	1,131	5%	60
POP - N of 40°10' N. lat. d/	133.5	95%	126.8	5%	6.7
Pacific whiting	TBA	100%	TBA	0%	TBA
Petrale sole a/	2,358.0	NA	2323.0	NA	35.0
Sablefish					
N of 36° N. lat.		See Table 1c of this subpart			
S of 36° N. lat.	1,434	42%	602	58%	832
Shortspine thornyhead					
N of 34°27' N. lat.	1,481	95%	1,407	5%	74
S of 34°27' N. lat.	355	NA	50	NA	305
Splitnose - S of 40°10 N. Lat.	1,598	95%	1,518	5%	80
Starry Flounder	1,513	50%	757	50%	757
Widow e/	1,411	91%	1,284	9%	127
Yelloweye rockfish a/	12.2	NA	1.0	NA	11.2
Yellowtail - N of 40°10 N. Lat.	3,677	88%	3,235	12%	441

- a/ Allocations decided through the biennial specification process.
- b/ 12.6 mt of the total trawl allocation of canary rockfish is allocated to the at-sea whiting fisheries, as follows: 5.2 mt for the mothership fishery, and 7.4 mt for the catcher/processor fishery.
- c/ 9 percent (25.3 mt) of the total trawl allocation for darkblotched rockfish is allocated to the whiting fisheries, as follows: 10.6 mt for the shorebased IFQ fishery, 6.1 mt for the mothership fishery, and 8.6 mt for the catcher/processor fishery. The tonnage calculated here for the whiting portion of the shorebased IFQ fishery contributes to the total shorebased trawl allocation, which is found at § 660.140(d)(1)(ii)(D).
- d/ 30 mt of the total trawl allocation for POP is allocated to the whiting fisheries, as follows: 12.6 mt for the shorebased IFQ fishery, 7.2 mt for the mothership fishery, and 10.2 mt for the catcher/processor fishery. The tonnage calculated here for the whiting portion of the shorebased IFQ fishery contributes to the total shorebased trawl allocation, which is found at § 660.140(d)(1)(ii)(D).
- e/ 500 mt of the total trawl allocation for widow rockfish is allocated to the whiting fisheries, as follows: 210 mt for the shorebased IFQ fishery, 120 mt for the mothership fishery, and 170 mt for the catcher/processor fishery. The tonnage calculated here for the whiting portion of the shorebased IFQ fishery contributes to the total shorebased trawl allocation, which is found at § 660.140(d)(1)(ii)(D).

[78 FR 592, Jan. 3, 2013]