

(3) On Anjou, smooth solid or smooth netlike russeting when the aggregate area exceeds one-third of the surface, and on other smooth-skinned varieties, 15 percent of the surface, except that, in addition, on Anjou and other smooth-skinned varieties, any amount of characteristic smooth russeting shall be permitted on that portion of the calyx end not visible for more than one-half inch along the contour of the pear, when it is placed calyx end down on a flat surface.

(4) On any of the following and other similar varieties, rough or thick russeting such as is characteristic of frost injury, when the aggregate area exceeds three-fourths inch in diameter. On any of these varieties any amount of characteristic russeting is permitted whether due to natural causes such as weather or stimulated by artificial means; leaf whips or light limbrubs which resemble and blend into russeted areas shall be considered as russet:

Bosc, Clairgeau, Comice, Easter Beurre, Flemish Beauty, Kieffer, P. Barry, Pound, Seckel, Sheldon, Winter Nelis, and other similar varieties.

(e) Any one of the following defects or any combination thereof, the seriousness of which exceeds the maximum allowed for any one defect, shall be considered as damage:

(1) Any limbrubs which are cracked, softened, or more than slightly depressed.

(2) Black discoloration caused by limbrubs which exceeds an aggregate area of three-eighths inch in diameter.

(3) Dark brown discoloration or excessive roughness caused by limbrubs which exceeds an aggregate area of one-half inch in diameter.

(4) Slightly rough, light colored discoloration caused by limbrubs which exceeds an aggregate area of three-fourths inch in diameter.

(5) Smooth, light colored discoloration caused by limbrubs which exceeds an aggregate area of 1 inch in diameter.<sup>2</sup>

(6) Hail marks or other similar depressions or scars which are not shallow or superficial, or where the injury affects an aggregate area of more than three-eighths inch in diameter.<sup>2</sup>

(7) Cork spot when more than one in number is visible externally or when the flesh is materially affected.

(8) Drought spot when more than one in number, or when the external injury exceeds an aggregate area of three-eighths inch in diameter, or when the appearance of the flesh is materially affected by corky tissue or brownish discoloration.<sup>2</sup>

(9) Sunburn or sprayburn where the skin is blistered, cracked, or shows any light tan or brownish color, or the shape of the pear is appreciably flattened, or the flesh is appreciably softened or changed in color, except that sprayburn of a russet character shall be considered under the definition of russeting.

(10) Insects: (i) More than two healed codling moth stings, or any insect sting which is over three thirty-seconds of an inch in diameter, or other insect stings affecting the appearance to an equal extent.<sup>2</sup>

(ii) Blister mite or canker worm injury which is not shallow or superficial, or where the injury affects an aggregate area of more than three-eighths inch in diameter.<sup>2</sup>

(11) Disease: (i) Scab spots which are black and which cover an aggregate area of more than one-fourth inch in diameter, except that scab spots of a russet character shall be considered under the definition of russeting.<sup>2</sup>

(ii) Sooty blotch which is thinly scattered over more than 5 percent of the surface, or dark, heavily concentrated spots which affect an area of more than three-eighths inch in diameter.<sup>2</sup>

#### § 51.1322 Seriously misshapen.

*Seriously misshapen* means that the pear is excessively flattened or elongated for the variety, or is constricted or deformed so it will not cut three fairly uniform good quarters, or is so badly misshapen that the appearance is seriously affected.

#### § 51.1323 Serious damage.

*Serious damage* means any injury or defect which seriously affects the appearance, or the edible or shipping quality.

(a) Russeting which in the aggregate exceeds the following shall be considered as serious damage:

(1) On all varieties, excessively rough russeting (russeting which shows "frogging" or slight cracking) when

the aggregate area exceeds three-fourths inch in diameter.<sup>2</sup>

(2) On all varieties, thick russetting such as is characteristic of frost injury, 15 percent of the surface.

(3) On Anjou, smooth solid or smooth netlike russetting when the aggregate area exceeds two-thirds of the surface, except that, in addition, any amount of characteristic smooth russetting shall be permitted on that portion of the calyx end not visible for more than one-half inch along the contour of the pear, when it is placed calyx end down on a flat surface. On Flemish Beauty smooth russetting shall be permitted on the entire surface.

(b) Any one of the following defects or combination thereof, the seriousness of which exceeds the maximum allowed for any one defect, shall be considered as serious damage:

(1) Limbrubs which are more than slightly cracked, or excessively rough limbrubs or dark brown or black discoloration caused by limbrubs which exceeds an aggregate area of three-fourths inch in diameter.<sup>2</sup>

(2) Other limbrubs which affect an aggregate area of more than one-tenth of the surface.

(3) Hail marks or other similar depressions or scars which affect an aggregate area of more than three-fourths inch in diameter, or which materially deform or disfigure the fruit.<sup>2</sup>

(4) Cork spot when more than two in number are visible externally or when the flesh is seriously affected.

(5) Drought spot when more than two in number, or where the external injury affects an aggregate area of more than three-fourths inch in diameter, or when the appearance of the flesh is seriously affected by corky tissue or brownish discoloration.<sup>2</sup>

(6) Sunburn or sprayburn where the skin is blistered, cracked or shows any brownish color, or where the shape of the pear is materially flattened, or the flesh is softened or materially changed in color, except that sprayburn of a russet character shall be considered under the definition of russetting.

(7) Insects: (i) Worm holes. More than three healed codling moth stings, of which not more than two may be over three thirty-seconds of an inch in di-

ameter, or other insect stings affecting the appearance to an equal extent.<sup>2</sup>

(ii) Blister mite or canker worm injury which affects an aggregate area of more than three-fourths inch in diameter or which materially deforms or disfigures the fruit.<sup>2</sup>

(8) Disease: (i) Scab spots which are black, and which cover an aggregate area of more than one-half inch in diameter, except that scab spots of a russet character shall be considered under the definition of russetting.<sup>2</sup>

(ii) Sooty blotch which is thinly scattered over more than 15 percent of the surface, or dark heavily concentrated spots which affect an area of more than three-fourths inch in diameter.<sup>2</sup>

**Subpart—United States Standards for Pears for Canning**

SOURCE: 35 FR 6957, May 1, 1970, unless otherwise noted. Redesignated at 42 FR 32514, June 27, 1977, and further redesignated at 46 FR 63203, Dec. 31, 1981.

**GENERAL**

**§ 51.1345 General.**

All percentages shall be calculated on the basis of weight.

**GRADES**

**§ 51.1346 U.S. No. 1.**

“U.S. No. 1” consists of pears of one variety which are mature, handpicked, firm, well formed, free from scald, hard end, black end, internal breakdown, decay, worms and worm holes, and from damage caused by broken skins, limbrubs, sprayburn, sunburn, scab, russetting, bruises, hail, frost, drought spot, disease, insects, mechanical or other means. Unless otherwise specified, the pears shall not be further advanced than yellowish green. Tree-ripened pears and pears grown from late blooms shall not be considered as meeting the requirements of this grade. (See §§ 51.1349 and 51.1350.)

**§ 51.1347 U.S. No. 2.**

“U.S. No. 2” consists of pears of one variety which are mature, handpicked, firm, not seriously deformed, free from scald, hard end, black end, internal breakdown, decay, worms and worm